



In vitro* Effect of Bio-agents for Control of Alternaria Blight of Indian Mustard Incited by *Alternaria brassicicola

**Ravi Regar ^{a*}, S. K. Goyal ^a, S. Godika ^a, Pinki Sharma ^a,
Dinesh Kumar Meena ^a, Pooja Kumawat ^a,
Vijay Kumar Kasotiya ^a and Suman Chopra ^a**

^a Department of Plant Pathology, S. K. N. College of Agriculture, Jobner - 303329, Jaipur (Rajasthan), India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2024/v36i95034>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/124011>

Original Research Article

Received: 18/07/2024

Accepted: 20/09/2024

Published: 26/09/2024

ABSTRACT

Indian mustard [*Brassica juncea* (L.) Czern and Coss.] is one of the most popular *rabi* oil seed crop in India. It's belonging to family Cruciferae (Brassicaceae). It is mostly cultivated *rabi*, crop of Northern as well as North eastern region of India. Indian mustard contributes about 85 per cent of total rapeseed-mustard production in India [1]. Crucial Indian mustard growing states are Rajasthan (50%), Uttar Pradesh (12.3%), Haryana (11.2%), Madhya Pradesh (9.8%), Gujrat (6.5%) and West Bengal (5.1%). Indian mustard is affected by many major diseases like Fungal, Bacterial, Viral, and also parasitic nematode. In mustard maximum losses in yield occurs due to microbial diseases. Among the major disease viz., Alternaria blight (*Alternaria brassicae*, *Alternaria brassicicola*, *Alternaria alternata*), White rust (*Albugo candida*), Downey mildew (*Peronospora parasitica*),

*Corresponding author: E-mail: raviregar4135@gmail.com;

Powdery mildew (*Erysiphe cruciferarum*), Bacterial blight (*Pseudomonas cannabina*), and Sclerotinia stem rot (*Sclerotinia sclerotiorum*). Alternaria blight of Indian mustard is one of the most important disease which reduce quantity and quality of seed yield. Alternaria blight disease is one of the damaging disease among the major diseases of mustard causing up to 70 per cent losses in yield with no proven source of resistance against the disease reported till date in any of the hosts. To control the losses caused by pathogens many farmers are using fungicides to obtain very good yield. The present study was carried out to recognize the effect of different bioagents on control against *Alternaria brassicicola* under *in-vitro* condition. Four antagonists viz., *Trichoderma viride*, *Trichoderma harzianum*, *Pseudomonas fluorescens* and *Bacillus subtilis*, checked the growth of *Alternaria brassicicola* at point of their contact on culture medium. Among these, maximum inhibition (78.05%) of *Alternaria brassicicola* was observed in *Trichoderma harzianum*.

Keywords: Indian mustard; bioagents; per cent inhibition; *Trichoderma harzianum*.

1. INTRODUCTION

“Indian mustard [*Brassica juncea* (L.) Czern and Coss.] is one of the most popular *rabi* oil seed crop in India. It's belonging to family Cruciferae (Brassicaceae). It is mostly cultivated *rabi*, crop of Northern as well as North eastern region of India. It was emanated in China and from there it was introduced to India. According to ancient scripture, it has been grown as early as 5000 BC” [2].

“Indian mustard contributes about 85 per cent of total rapeseed-mustard production in India” [1]

“Mustard seed oil is popularly used for edible purpose, but also it used as industrial lubricant. Mustard cake is used as organic manure and as a protein course in animal feeds” [3].

Indian mustard is affected by many major diseases like Fungal, Bacterial, Viral, and also parasitic nematode. In mustard maximum losses in yield occurs due to microbial diseases. Among the major disease viz., Alternaria blight (*Alternaria brassicae*, *Alternaria brassicicola*, *Alternaria alternata*), White rust (*Albugo candida*), Downey mildew (*Peronospora parasitica*), Powdery mildew (*Erysiphe cruciferarum*), Bacterial blight (*Pseudomonas cannabina*), and Sclerotinia stem rot (*Sclerotinia sclerotiorum*).

“Alternaria blight of Indian mustard is one of the most important disease which reduce quantity and quality of seed yield. In Varuna cultivar under unprotected conditions 20.8 per cent seed yield loss” observed by Prasad et al. [4].

According to Chadar et al. [5] 23.9 to 62 per cent disease intensity of Alternaria leaf blight of mustard which occurred extensively in Jhansi region of India.

“Symptoms of disease appear on leaves and stems and adult plants. Due to Alternaria disease dark brown spots on the leaves and siliquae reduce the photosynthetic capacity and induce immature ripening, which causes decreased the quality seed production in both vegetable and brassicas” [2].

“Chemicals are not only costly but also they are creating problems on the environment, human health in all areas of the world” [6]. Now a day bioagents also used for control of many fungal diseases because those are not hazardous to crop as well as humans.

The Present studied was carried out to find the best botanical bioagent on control against *Alternaria brassicicola* causing Alternaria blight of Indian mustard.

2. MATERIALS AND METHODS

2.1 Isolation of Fungus Cultures

“The diseased plants were collected from the mustard grown in the field. The Alternaria blight affected leaf and stem portion was cut into small pieces and surface sterilized with 0.1% sodium hypochlorite solution then afterward three times thoroughly washed with water” [7]. Potato dextrose agar (PDA) media were prepared and sterilized in an autoclave at 120°C, 15 lbs pressure for 20 minutes. The media were poured into Petri plates and allowed to cool for sometimes. Then, the leaf pieces were inoculated into the media and kept for incubation. After seven to eight days the fungal tips were transferred to PDA slants in order to obtain pure cultures. The isolates were confirmed as *Alternaria* [8] by microscopic observation of fungal culture.

List 1. Experimental details

Design: C.R.D. (Completely Randomized Design)	
Replication: Four	
Treatments: Five	
S. No.	Bioagents
1	<i>Trichoderma viride</i>
2	<i>Trichoderma harzianum</i>
3	<i>Pseudomonas fluorescens</i>
4	<i>Bacillus subtilis</i>
5	Control

2.2 In vitro Management of Disease through Bioagents

“Two fungal and two bacterial bioagents were evaluated *in vitro* against *Alternaria brassicicola* applying Dual Culture Technique” [9]. “Seven days old cultures of the test bioagents and test pathogen (*Alternaria brassicicola*) grown on PDA were used for the study. Two 5 mm culture discs, one each of the test pathogen and test bioagent were cut out with sterilized cork borer and placed at equidistance, exactly opposite to each other on autoclaved and solidified PDA medium in Petri plates and three plates were incubated at 28 ± 2 °C PDA plates inoculated alone with pure culture disc (5 mm) of the test pathogen were maintained as untreated control” [7].

The extent antagonistic activity by bioagent was recorded after full growth covers in the control plate.

The per cent mycelial inhibition of pathogen was calculated using formula of Bliss [10].

$$I = \frac{(C-T)}{C} \times 100$$

Where, I = Per cent mycelial inhibition

C = Growth of fungal pathogen in Control (mm)
T = Growth of fungal pathogen in dual culture plate (mm)

3. RESULTS AND DISCUSSION

The results of fungal mycelial growth and *Alternaria brassicicola* suppression with two fungal and two bacterial bioagents are shown in (Table 1 and Plate 1).

Results showed that all four bioagents evaluated exhibited antifungal activity against *Alternaria brassicicola* and significantly inhibited pathogen growth over control. *Trichoderma harzianum* was superior to *Trichoderma viride*, *Bacillus subtilis* and *Pseudomonas fluorescens* which produced 78.05 per cent inhibition of mycelial growth of pathogen in dual culture plates (Table 1 and Fig. 1) These findings were in accordance with Zafar et al. [11] and Selim [12]. Sivapalan [13] observed that *Gliocladium roseum* and *Trichoderma harzianum* reduced leaf infection of *Alternaria brassicicola* on broccoli. Prasad and Kulshrestha [14] observed that *Pseudomonas fluorescens* showed lowest incidence of blight infected seedling of sunflower followed by *Bacillus subtilis* [15].

Table 1. In vitro evaluation of bioagents against *Alternaria brassicicola*

S. No.	Bioagents	Mycelial Growth (mm)	Per cent inhibition of mycelial growth
1	<i>Trichoderma viride</i>	28.50	68.33 (55.73)
2	<i>Trichoderma harzianum</i>	19.75	78.05 (62.04)
3	<i>Pseudomonas fluorescens</i>	54.75	39.16 (38.72)
4	<i>Bacillus subtilis</i>	52.50	41.67 (40.18)
5	Control	90	0.00
	SEm±	0.671	
	CD (p=0.05)	2.091	
	CV	2.363	




Fig. 1. *In vitro* evaluation of bioagents against *Alternaria brassicicola*

**Mean of four replications*

***Figures in the parenthesis are square root transformed values*




Plate 1. *In vitro* evaluation of bioagents against *Alternaria brassicicola*

4. CONCLUSION

Among four bioagents, *Trichoderma harzianum* was found most effective against *Alternaria brassicicola* *In vitro* condition. The present study was carried out to recognize the effect of different bioagents on control against *Alternaria brassicicola* under *in-vitro* condition. Four antagonists viz., *Trichoderma viride*, *Trichoderma harzianum*, *Pseudomonas fluorescens* and *Bacillus subtilis*, checked the growth of *Alternaria brassicicola* at point of their contact on culture medium.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

ACKNOWLEDGEMENT

I would like to acknowledge to the Head of the department of Plant Pathology, College of Agriculture, Jobner, Jaipur, for providing me necessary facilities and motivated for conducting the research work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kumar A, Chauhan JS. Strategies and future thrust areas of rape seed mustard research in India. *Indian Journal Agriculture Science*. 2005;75:621-635.
2. Kumar D, Maurya N, Bharti YK, Kumar A, Kumar K, Srivastava K, Chand G, Kushwaha C, Singh SK, Mishra RK, Kumar A. *Alternaria* blight of oilseed Brassicas: A comprehensive review. *African Journal of Microbiology*. 2014;8(30):2816-2829.
3. Bhowmik TP. *Oilseed Brassicas constraints and their management*. CBS Publishers and Distributors, New Delhi. 2003;1-2.
4. Prasad R, Saxena D, Chandr S. Yield losses by *alternaria* blight in promising genotypes of Indian mustard. *Indian Phytopathology*. 2003;56(2):205-206.
5. Chadar LK, Singh RP, Singh RK, Yadav RR, Mishra MK, Pratap N, Vishnoi RK. Studies on *alternaria* blight of rapeseed-mustard (*Brassica juncea* L.) caused by *alternaria brassicae* (Berk.) Sacc. and its integrated management. *Plant Archives*. 2016;16(2):897-901.
6. Rahmatzai N, Zaitoun AA, Madkour MH, Ahmady A, Hazim Z, Mousa MA. *In-vitro* and *In-vivo* antifungal activity of botanical oils against *Alternaria solani* causing early blight of tomato. *International Journal of Biosciences*. 2017;10(1):91-99.
7. Ravi Regar, Bireswar Sinha, Veena Khot, Bhuvaneswari, Biram Singh Gurjar. *In vitro* Cultural characteristics and effect of botanical (Garlic) for control of major fungal diseases of potato in Manipur. *Int. J. Curr. Microbiol. App. Sci*. 2021;10(03):733-739.
8. Van Bruggen AH. Sweet potato stem blight caused by *Alternaria* spp.: A new disease in Ethiopia. *Netherlands Journal of Plant Pathology*. 1984;90(4):155-164.
9. Dennis C, Webster J. Antagonistic properties of species-groups of *Trichoderma*: I. Production of non-volatile antibiotics. *Transactions of the British Mycological Society*. 1971;57(1):25-33.
10. Bliss CL. The method of probits. *Science*. 1934;79:38
11. Zafar MN, Wahid A, Ghauri MA, Zubair M, Mumtaz MW, Anwar F, Danish M. *Trichoderma harzianum*: a green sorbent for Pb (II) uptake from aqueous solutions. *Toxicological & Environmental Chemistry*. 2013;95(3):422-433.
12. Selim ME. Effectiveness of *Trichoderma* biotic applications in regulating the related defence genes affecting tomato early blight disease. *Journal of Plant Pathology and Microbiology*. 2015;6(311):2
13. Sivapalan A. Fungi associated with broccoli seed and evaluation of fungal antagonists and fungicide for the control of seed borne *Alternaria brassicicola*. *Seed Science and Technology*. 1993;21(1):237-245
14. Prasad Kulshrestha. Efficacy of some bacterial antagonists against seedling blight of sunflower caused by *Alternaria helianthi*. *Plant Disease Resistance*. 2002;17(1):60-65.

15. Ahmad F, Raziq F, Ullah N, Khan H, Din N. *In vitro* and *In vivo* bio-assay of phytobiocidal effect of plant extracts on *Alternaria solani* causing agent of early blight disease in tomato. Archives of Phytopathology and Plant Protection. 2017;50(11-12):568-583.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/124011>