



Resistance Pattern of Nitrofurantoin of Uropathogens in Different Age Groups at Dr. Lal Path Labs, National Reference Laboratory, Rohini, Delhi

Puneeta Singh¹, Shalabh Malik^{1*} and Vandana Lal²

¹*Department of Microbiology and Serology, Dr. Lal Path Labs, National Reference Laboratory, Rohini, Delhi, India.*

²*Dr. Lal Path Labs, National Reference Laboratory, Rohini, Delhi, India.*

Authors' contributions

This work was carried out in collaboration among all authors. Author SM designed the study, wrote the protocol, analyzing the protocol and review of manuscript. Author PS wrote the first draft of the manuscript, performed the statistical analysis and managed the literature searches. Author VL managed the analyses of the study and infrastructure support. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJMAH/2019/16i130134

Editor(s):

(1) Professor, Janvier Gasana, Department of Environmental and Occupational Health, EO Epidemiology and EO Medicine, Robert Stempel College of Public Health and Social Work, Florida International University, USA.

(2) Dr. Nicolas Padilla – Raygoza, Professor, Department of Nursing and Obstetrics, Division of Health Sciences and Engineering, Campus Celaya-Salvatierra, University of Guanajuato, Mexico.

Reviewers:

(1) Ramachandra Reddy, Deemed University, India.

(2) Teresita Sainz-Espuñes, UAM, Mexico.

Complete Peer review History: <http://www.sdiarticle3.com/review-history/50908>

Original Research Article

Received 11 June 2019
Accepted 17 August 2019
Published 24 August 2019

ABSTRACT

Aims and Objectives: This study was undertaken to highlight the resistance pattern of Nitrofurantoin among 5162 UTIs causing isolates at Microbiology Department of Dr. Lal Path Labs between April to June 2019.

Materials and Methods: This retrospective study was performed in Department of Microbiology at Dr. Lal Path Labs, Delhi during period April to June 2019. Standard loopful midstream urine samples collected in a sterile container were inoculated on UTI Chromagar and incubated overnight at 37°C and demonstrating significance colony count of $\geq 10^5$ CFU/ml. Common Enterobacteriaceae

*Corresponding author: E-mail: Shalabh.malik@lalpathlabs.com;

group of isolates and Gram positive isolates identified by MALDI TOF-MS (Bruker, Daltonics) were included in this study. To determine Nitrofurantoin resistance by VITEK-AST (Biomerieux) system on 280/P628 antibiotic susceptibility testing cards respectively as per as CLSI M100-S-29. *Proteus spp*, *Pseudomonas spp.*, *Acinetobacter spp.*, of complicated UTI isolates were excluded from this study.

Results: Of the 29485 urine samples tested during April to June 2019, 5162 (17.5%) were culture positive. Out of the 5162 positive isolates 2856 (55.3%) were isolated from female patients and 2306 (44.7%) from male patients. The most common bacterial isolates were members of Enterobacteriaceae 4728 (91.5%) and Gram positive were 434 (8.5%). The most predominant age group infected with uropathogens were elderly adults ≥ 50 years of age constituted 58.1% of the UTIs culture positive cases. *Klebsiella pneumoniae* demonstrated highest resistance of 92.3% whereas *Staphylococcus aureus* demonstrated the least resistance of 8.3%.

Discussion and Conclusion: The alarming substantial resistance to Nitrofurantoin in decreasing order has been noticed in Enterobacteriaceae i.e., *Klebsiella pneumonia* (92.3%), *Enterobacter spp.*(58.2%), *Citrobacter spp* (42.6%) and Gram positive *Enterococcus spp.* (45.6%) which is across all age groups. Most susceptible to Nitrofurantoin were *Escherichia coli* (69.9%) in Gram negative and *Staphylococcus aureus* (91.7%) in Gram positive. This finding emphasis the need of robust restriction of Nitrofurantoin antibiotic policy and usage to uncomplicated UTIs caused by *Escherichia coli* and *Staphylococcus aureus*.

Keywords: Nitrofurantoin; uropathogens; UTIs (Urinary Tract Infection); Enterobacteriaceae; gram positive.

1. INTRODUCTION

Urinary UTIs, is one of the most frequent infection in mankind, and are still among the most common bacterial infections in the world. It is estimated to affect 150 million people each year world wide. [1] Nitrofurantoin is a broad spectrum, cheap and best bactericidal antibiotic which is used for treating uncomplicated UTIs and nosocomial lower UTIs, that is an antibiotic for affects both Gram -ve and Gram+ve bacteria including *Escherichia coli*, *Klebsiella pneumoniae*, *Citrobacter*, *Enterobacter*, *Enterococcus*, *Staphylococcus aureus*, ESBL producing strains, also active against VRE and VSE [1,2]. Nitrofurantoin is active against most common uropathogens but most *Proteus species*, *Serratia marcescens* and *Pseudomonas aeruginosa* are naturally resistant [2].

Incidence, prevalence and antibiogram of adults and pediatric UTIs differ from country to country and within same country between different geographical areas and also in different age groups [3]. The alarming rise of resistant to Nitrofurantoin is a matter of concern about the use of Nitrofurantoin limitations in Indian scenario.

It was observed that increasing of MIC of Nitrofurantoin in pediatric and adults UTIs has not been reported from this part of our country that is northern India, Delhi. In this study, the

objective was to investigate the Cumulative MIC of Nitrofurantoin for different types of uropathogens. On the basis of our findings Nitrofurantoin should no longer be recommended for initial empirical therapies for *Klebsiella pneumoniae*, *Enterobacter*, *Enterococcus*, noscomial *E. Coli*. Hence this study was undertaken to determine the cumulative interpretation and MIC of Nitrofurantoin of pediatric and adults UTIs.

2. METHODS

This retrospective study was performed in Department of microbiology at Dr Lal Path Labs, Delhi for a period April to June 2019. A total of 29,485 midstream urine samples were submitted to microbiology department of *Dr Lal Path Labs* for processing. According to the standard microbiological techniques with standard (10 μ l) loopful urine was inoculated on UTI Chrom agar and incubated overnight at 37°C under aerobic conditions. Based on Cfu/ml, the cultures were classified as negative, insignificant, Significant and contamination as per standard recommendations. Significant growth was determined as $>10^5$ colony forming units CFU/ml of midstream urine, $>10^2$ CFU/ml of a catheter specimen and any no. of colonies from a suprapubic sample. More than two types of bacteria on culture were excluded from this study.

Common Enterobacteriaceae group of isolates *Escherichia coli*, *Klebsiella pneumoniae*, *Citrobacter*, *Enterobacter*, and Gram positive isolates of *Enterococcus*, *Staphylococcus aureus* identified by MALDI TOF-MS (Bruker, Daltonics) were included in this study. VITEK-2 (Biomerieux) system was employed for the antibiotic susceptibility testing of isolates from the pure culture of isolated colonies of the uropathogens on UTI Chrom agar, the Gram negative and Gram positive bacteria were inoculated on to N280/P628 cards respectively.

CLSI M100-S-29 [4] interpretive criteria for Enterobacteriaceae and Gram positive were utilized for Nitrofurantoin. For Enterobacteriaceae, *Enterococcus*, *Staphylococcus aureus* breakpoints were $\leq 32\mu\text{g/ml}$ (susceptible), $64\mu\text{g/ml}$ (intermediate) and $\geq 128\mu\text{g/ml}$ (resistant). *Proteus spp.*, *Pseudomonas spp.*, *Acinetobacter spp.*, of complicated UTI isolates were excluded from this study.

2.1 Statistical Analysis

For the evaluation of the study data Myla (Bio Merieux, India Pvt. Ltd).

Statistical analysis program was used.

3. RESULTS

We assessed the activity of Nitrofurantoin against 5162 (17.5%) that comprises Enterobacteriaceae and Gram positive UTIs uropathogens that were collected during the study period, they consisted of *Escherichia coli* (71.5%), followed by *Klebsiella pneumoniae* (17.8%), *Citrobacter spp.* (0.7%), *Enterobacter spp.* (1.5%), *Enterococcus spp.* (7.7%), *Staphylococcus aureus* (0.7%) (Fig. 1). Out of the 5162 positive isolates 2856 (55.3%) were isolated from female patients and 2306 (44.7%) from male patients. 6.1% of isolates of them belonging to the age group of 0-12 years and rest were 93.9% of 13-95 years. In our study the most predominant age group infected with uropathogens were elderly adults ≥ 50 years (58.1%) followed by adults (20%), young adults (15.8%) and children (6.1%) (Table 1). The prevalence of isolates among Enterobacteriaceae and Gram positive were stratified by age group, the most frequently identified bacteria in pediatric age group were *Escherichia coli* (76.1%) (Fig. 2). Antibiotic resistance to Nitrofurantoin was elevated across all age groups for Enterobacteriaceae and Gram

positive bacterial species, but it was especially high among isolates of *Klebsiella pneumoniae* (92.3%), *Enterobacter* (58.2%) and *Enterococcus* (45.6%). Conversely *Escherichia coli* (30.1%) and *Staphylococcus aureus* (8.3%) resistance to Nitrofurantoin were low (Table 2). The prevalence of resistance increased by age group for several uropathogens for example, *Citrobacter*, *Staphylococcus aureus* resistant to Nitrofurantoin were relatively low among isolates from pediatric age group (Fig. 3).

Table 1. Distribution of uropathogens in different age groups

Age groups	Total number of uropathogens	% of uropathogens
0-12	314	6.1
13-30	814	15.8
31-50	1035	20
>51-95	2999	58.1

The cumulative interpretation of Nitrofurantoin drug in different age groups shown dramatically changes we found that resistance of Nitrofurantoin were not related to age groups all uropathogens consistently increasing in all age groups. Only *Staphylococcus aureus* shown sensitive against Nitrofurantoin and most predominant age group were ≥ 51 year of age (Fig. 3).

And this study describes for the first time cumulative MIC interpretation of Nitrofurantoin resistance patterns among Enterobacteriaceae and Gram positive isolates with help of Myla statistical analysis (Biomerieux, India) which causes complicated UTIs such as *Klebsiella pneumoniae*, *Enterobacter* and *Enterococcus* is increasing in community acquired UTIs in Delhi. Total 3691(71.5%) *Escherichia coli* isolates tested against Nitrofurantoin, 45% of *Escherichia coli* isolates was having MIC $\leq 16\mu\text{g/ml}$ and 70% of isolates tested was having MIC $\leq 32\mu\text{g/ml}$ (Table 3). Out of 921(17.8%) tested isolates of *Klebsiella pneumoniae* only 3% isolates having MIC $\leq 16\mu\text{g/ml}$ and 8% of isolates was having MIC $\leq 32\mu\text{g/ml}$. Nitrofurantoin activity (MIC_{50/90} 128/512) against *Klebsiella pneumoniae* demonstrated that 50% of isolate were within $128\mu\text{g/ml}$ MIC and 90% isolates were within $512\mu\text{g/ml}$, *Klebsiella pneumoniae* were recorded high resistance rate (92.3%) in this study (Table 3). Second highest resistance recorded to Nitrofurantoin in *Enterobacter spp.* (58.2%) in Delhi.

Table 2. Percentage of cumulative interpretation of nitrofurantoin resistance in all age groups

<i>Escherichia coli</i>	<i>Klebsiella pneumoniae</i>	<i>Citrobacter spp.</i>	<i>Enterobacter spp.</i>	<i>Enterococcus spp.</i>	<i>Staphylococcus aureus.</i>
30.1%	92.3%	42.6%	58.2%	45.6%	8.3%

Distribution of uropathogens among culture positive samples in all age groups during April to June 2019.




Fig. 1. Distribution of uropathogens among culture positive samples in all age groups during April to June 2019

Table 3. Percentage of cumulative MIC interpretation and antimicrobial activity of nitrofurantoin against uropathogens from all age groups during April to June 2019

Uropathogens	MIC ($\mu\text{g/ml}$)/ cumulative%							MIC ₅₀	MIC ₉₀	%S	%R
	16	32	64	128	256	512					
<i>Escherichia coli</i>	45	70	90	96	99	100	32	128	69.9	30.1	
<i>Klebsiella pneumoniae</i>	3	8	32	58	78	100	128	512	7.7	92.3	
<i>Citrobacter spp.</i>	13	57	76	89	95	100	64	256	57.4	42.6	
<i>Enterobacter spp.</i>	8	42	76	100	--	--	64	128	41.8	58.2	
<i>Enterococcus spp.</i>	28	54	63	82	93	100	64	256	54.4	45.6	
<i>Staphylococcus aureus</i>	50	92	90	90	100	---	≤ 16	64	91.7	8.3	

-- -Not tested

Distribution of uropathogens in different age groups during April to June 2019.




Fig. 2. Age specific distribution of uropathogens during April to June 2019

Percentage of resistance patterns of Nitrofurantoin in uropathogens on the basis of cumulative interpretation in different age groups during April to June 2019.




Fig. 3. Percentage of cumulative interpretation of nitrofurantoin resistance patterns of age specific distribution of uropathogens during April to June 2019

The distribution of Nitrofurantoin MIC values against resistant patterns of other uropathogens followed in (Table 3).

4. DISCUSSION

Many countries as well as different part of India reported Nitrofurantoin as first line of antibiotic for treatment and prophylaxis of acute lower UTIs

and noscomial infection of UTIs [1,5,6,7,8]. Our study describes the distribution and antibiotic resistance of Nitrofurantoin based on Cumulative interpretation and MIC across all age groups.

Comparing the age groups most commonly affected by UTIs with different uropathogens in our study was the elderly group aged ≥ 50 age

and least affected (0-12) years of age which is similar to other studies [7,3].

This study highlighted potential and the limitation of this agent in the era of antibiotic resistance especially in Delhi, India. *Escherichia coli*, *Klebsiella pneumoniae*, *Citrobacter*, *Enterobacter* and *Enterococcus* isolates are reported to be the most common organisms causing UTIs in not only noscomial infections but community acquired infections.

In accordance with the several global and national reports our study revealed *Escherichia coli* (71.5%) as the most predominantly isolated uropathogen associated with UTIs in all age groups [1-12,13,8]. Throughout the entire study, 30.1% of *Escherichia coli* isolates showed resistant against Nitrofurantoin. Similar finding were also reported by several authors [7,8,3].

The present study of cumulative MIC of Nitrofurantoin resistance have reported high level of resistance on *Klebsiella pneumoniae* (92.3%), which is in agreement with the findings of few studies from India and Taiwan quoted high resistance against Nitrofurantoin (>75%) in *Klebsiella pneumoniae* [14,15,16] Interesting thing is that the world seem Nitrofurantoin is sensitive in world for treatment and prophylaxis of *Klebsiella pneumoniae* in lower UTIs and noscomial infection [1-13,8,3].

To best of our knowledge 58.2% and 42.6% isolates of *Enterobacter*, *Citrobacter*, respectively that causes complicated UTIs having resistant to Nitrofurantoin. This finding was in contrast with previously performed studies in which *Citrobacter spp.* were reported sensitive to Nitrofurantoin [7,8,3]. Among the Gram negative organisms isolated in our study *Citrobacter spp.* (42.6%) and *Enterobacter spp.* (58.2%) had a high level of resistance to Nitrofurantoin, this is in consistence with findings of other studies [17,14].

On concordance to the finding of various other previous studies which documented among the gram positive organisms *Enterococcus spp.* (45.6%) showed very high level of resistance to nitrofurantoin [18] This finding was in contrast with previously performed studies in which *Enterococcus species* were reported sensitive to Nitrofurantoin [7,8,3].

Among the gram positive organisms isolated in our study *Staphylococcus aureus* had a very low

level of resistance (8.3%) to Nitrofurantoin used in this study this is similar with other studies [3].

Study of all uropathogens indicate that resistance to Nitrofurantoin is on rise and treatment of UTIs is becoming more difficult with time more over there are considerable regional and geographic differences in the susceptibility pattern of uropathogens is required and choose the appropriate empiric therapy of Nitrofurantoin for UTIs in children and adults.

To our knowledge, this is the first study that highlights MIC of Nitrofurantoin for *Escherichia coli*, *Klebsiella pneumoniae*, *Citrobacter spp.*, *Enterobacter spp.*, *Enterococcus spp.* and *Staphylococcus aureus* MIC₅₀ (concentration that inhibited 50% of isolates) was 32, 128, 64, 64, 16µg/ml and MIC₉₀ (concentration that inhibited 90% of isolates) was 128, 512, 256, 128, 256, 64 µg/ml respectively our results clearly demonstrated that Nitrofurantoin remains available suitable option for community acquired UTIs from *Escherichia coli* and *Staphylococcus aureus* in Delhi. This is in similar to other studies [8,19,20].

5. CONCLUSION

To conclude that UTIs varies with age groups therefore, extensive evaluation among interpretation by cumulative MIC of Nitrofurantoin increases with increasing age groups. Emergence of increasing MIC of Nitrofurantoin to *Klebsiella pneumoniae*, *Enterobacter spp.*, *Enterococcus spp.*, *Citrobacter spp.* has become the concern for policy makers and a urgent need of strict antibiotics prescription policy in our country. Judicious selection of antibiotics as per organisms recommendation by CLSI M-100. S-29 is the need of hour. Further Nitrofurantoin should be restricted to complicated and noncomplicated UTI by *Klebsiella pneumoniae* only.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

ACKNOWLEDGEMENTS

We thank staff of the Microbiology Department and Dr. Reena Nakra, Lab director, National

Reference Laboratory, Dr. Lal Path Labs, Rohini, Delhi.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Alqasim A, Jaffal AA, and Abdullah AA, et al. Prevalence of multidrug resistance and extended-spectrum β -lactamase carriage of clinical uropathogenic *Escherichia coli* isolates in Riyadh, Saudi Arabia. International Journal of Microbiology; 2018. Available: <https://doi.org/10.1155/2018/3026851>
2. Cunha BA, Schoch EP, Hage RJ et al., Mayo Clin Proc. 2011;86(12):1243-1244.
3. Singh RM, Devi MU, Singh KL, et al. Evaluation of nitrofurantoin activity against the urinary isolates in the current scenario of antimicrobial resistance. Ann Trop Med Public Health. 2015;8(6):280-285.
4. Clinical and Laboratory Standards Institute. Performance standards for antimicrobial disk susceptibility tests. Wayne Pa: Clinical and Laboratory Standards Institute: M100-S29; 2019.
5. Keyhan H, Sedighi S, Mashayekhi B, et al. Community acquired urinary tract infections, etiological organisms and antibiotics susceptibility patterns, Nephro-Urol Mon. 2017;9(5):e62146. DOI: 10.5812/NUMONTHLY.62146
6. Ekwealor PA, Malachy CU, Ezeobi I, Amalukwe G, Ugwu BC, Okezie U, Stanley C, Esimone C. Antimicrobial evaluation of bacterial isolates from urine specimen of patients with complaints of urinary tract infections in Awka, Nigeria. J Int J Microbiol. 2016;9740273.
7. Patwardhan V, Kumar D, Goel V, Singh S. Changing prevalence and antibiotic drug resistance pattern of pathogens seen in community-acquired pediatric urinary tract infections at a tertiary care hospital of North India. Journal of Laboratory Physicians. 2017;9(4):264-268.
8. Shakti L, Veeraraghavan B. Advantage and limitations of nitrofurantoin in multidrug resistant Indian scenario. Indian J Med Microbiol. 2015;33(4):477-81.
9. Maria Jose Munoz-Davila. Review, Antibiotics. 2014;3:39-48.
10. Gunduz S, Altun HU. Antibiotic resistance patterns of urinary tract pathogens in Turkish children. Global Health Research and Policy 2018; 3:10. Available: <https://doi.org/10.1186/s41256-018-0063-1>.
11. Huttner A, Verhaegh EM, Harbarth S, et al., Nitrofurantoin revisited: A systematic review and meta analysis of controlled trials, J Antimicrob chemother. 2015;70: 2456-2464.
12. Sanchez GV, Babiker A, Master RN, et al. Antibiotic resistance among urinary isolates from female outpatients in the United States in 2003 and 2012. Antimicrobial agents and chemotherapy; 2016. DOI: 10.1128/AAC.02897-15.
13. Nahar A, Hasnat S, Akhter H, et al. Evaluation of antimicrobial resistance pattern of uropathogens in a tertiary care hospital in Dhaka city, Bangladesh. South East Asia Journal of Public Health. 2017;7 (2):12-18,
14. Shailaja ST, Kumar MA. Antimicrobial resistance among uropathogenic bacteria in Rural Kerala, India. Int. J. Curr. Microbiol. App. Sci. 2017;6(9):2287-2296. DOI: <https://doi.org/10.20546/ijcmas.2017.609.2>
15. Mishra MP, Debata NK, and Pathy RN. Surveillance of multidrug resistant uropathogenic bacteria in hospitalized patients in India. Asian. Pac. J. Trop. Biomed. 2013;3(4):315-324.
16. Liu HY, Lin HC, Lin YC, Yu SH, Wu WH, Lee YJ. Antimicrobial susceptibilities of urinary extended spectrum beta lactamase-producing *Escherichia coli* and *Klebsiella pneumoniae* to fosfomycin and nitrofurantoin in a teaching hospital in Taiwan. J Microbiol. Immunol Infect. 2011; 44:364-8.
17. Patel HB, Soni ST, Bhagyalaxmi A and Patel NM. Causative agents of urinary tract infections and their antimicrobial susceptibility patterns at a referral center in Western India: An audit to help clinicians prevent antibiotic misuse. J Family Med Prim Care. Journal of Family Medicine and Primary care. 2019;8(1):154-159.
18. Benachinmardi K, Padmavathy M, Malini J, Navneeth BV. Microbiological profile and antibiogram of uropathogen in pediatric age group. Int. J. Health. Allied Science. 2015;4:61-64.

19. Komp Lindgren P, Klockars O, Malmberg C, Cars O. Pharmacodynamic studies of nitrofurantoin against common uropathogens. J. Antimicrob chemother. 2015;70(4):1076-82.
20. John Osei Sekyere. Genomic insights into nitrofurantoin resistance mechanism and epidemiology in clinical enterobacteriaceae. Future Sci OA. 2018; 4(5):FSO293.

© 2019 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<http://www.sdiarticle3.com/review-history/50908>