



# Relationship between Premature Rupture of the Membrane and Cesarean Delivery: Case from Jakarta, Indonesia

Jumaini Andriana Sihombing <sup>a\*</sup>, Muhammad Miqbel <sup>a</sup>  
and Batara Imanuel Sirait <sup>b</sup>

<sup>a</sup> Department of Anatomy, Faculty of Medicine, Universitas Kristen Indonesia, Jakarta, Indonesia.

<sup>b</sup> Department of Obstetrics and Gynaecology, Medical Faculty, Universitas Kristen Indonesia, Jakarta, Indonesia.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJRID/2023/v12i4253

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/97587>

Original Research Article

Received: 24/01/2023

Accepted: 26/03/2023

Published: 11/04/2023

## ABSTRACT

Premature rupture of membranes (PROM) is the spontaneous rupture of the membranes before the onset of signs of labor. It can occur both before term (<37 weeks) and after term (> 37 weeks). *Caesarean delivery* is a surgical technique to deliver the fetus through an incision in the abdominal wall and uterine wall. This procedure is performed to minimize all complications from premature rupture of membranes. This study aims to determine the relationship between premature rupture of membranes and delivery of *sectio caesarea* at the UKI General Hospital in the 2019 period by using a descriptive content analysis method (analysis of medical records documents) with a cross-sectional approach and obtained 65 samples who experienced premature rupture of membranes. The results showed that 53 samples (81.55%) experienced premature rupture of membranes and

\*Corresponding author: E-mail: jumaini.andriana@uki.ac.id;

performed *sectio caesarea*, while 12 samples (18.45%) had premature rupture of membranes but did not perform *sectio caesarea* with p-value = 0.001 ( $p < 0.05$ ) from the results of the chi-square statistical test. From the results of the analysis, it was found that there was a relationship between premature rupture of membranes and delivery of *cesarean section* at the UKI General Hospital for the 2019 period. It is suggested that patients avoid the factors that can cause premature rupture of the membranes, which have been told by the doctor or health worker and the patient's family to support the patient to remind them to avoid factors that can cause premature rupture of the membranes.

**Keywords:** *Premature rupture of membranes; cesarean section.*

## 1. INTRODUCTION

Premature or premature rupture of membranes is the spontaneous rupture of the amniotic membranes before signs of labor occur. This condition can occur either before term (<37 weeks) or after term (>37 weeks) [1]. The amniotic membranes function to produce amniotic fluid and protect the fetus from various kinds of infections [2]. Under normal circumstances, the membranes will rupture during labor. According to the World Health Organization, in 2014, the incidence of premature rupture of membranes or premature rupture of membranes ranged from 5-10% of all births. Preterm premature rupture of membranes occurs in 1% of all pregnancies, and 70% of premature rupture occurs at term. In 30% of cases, premature rupture of membranes is the cause of preterm birth [3].

The incidence of premature rupture of membranes in Indonesia ranges from 4.5% - 6% of all pregnancies. In contrast, in the United States, premature rupture of membranes occurs in 120,000 pregnancies per year and is associated with a high risk to maternal, fetal, and neonatal safety [4]. Various studies in India explain that the incidence is 7-12% for premature rupture of membranes which 60-70% occurs in the long term [5]. The incidence of premature rupture of membranes in several hospitals in Indonesia is quite varied, including Sardjito Hospital at 5.3%, Hasan Sadikin Hospital at 5.05%, Cipto Mangunkusumo Hospital at 11.22%, Pringadi Hospital at 2.27%, and Kariadi Hospital which is equal to 5.10% [6]. Premature rupture of membranes has various complications for the mother and fetus that cause morbidity and mortality. Maternal complications include chorioamnionitis and fetal complications such as fetal distress, premature birth, and cord compression, to neonatal complications such as neonatal asphyxia and sepsis. This complication requires labor conditions to be ended quickly to

save the mother and fetus. Sectio Caesarea is a labor procedure performed to minimize all complications from premature rupture of membranes [7].

Sectio Caesarea is a surgical technique to deliver a fetus through an incision in the abdominal wall (laparotomy) and uterine wall (hysterectomy). Sectio Caesarea is one of the ten most commonly performed major surgeries. This medical procedure is needed to assist deliveries that cannot be carried out normally due to maternal health problems or fetal conditions [8].

According to Travassolo in 2010 stated that premature rupture of membranes with oligohydramnios is related to the duration of rupture of membranes, high rates of sectio caesarea, and low Apgar scores at birth.

In Indonesia, the number of caesarean section deliveries continues to increase. According to the Indonesian Demographic and Health Survey (IDHS) data, there was an increase in caesarean sections in Indonesia from 1991 to 2007, around 1.3-6.8 percent [9]. Meanwhile, the results of the Riskesdas report (basic health research) in 2013 found that 9.8% of deliveries were carried out by sectio caesarea in Indonesia, 2.3% of which were due to complications in pregnancy [10].

Research conducted by Andayasari et al. in 2014 stated that sectio caesarea deliveries in one government hospital and one private hospital in DKI Jakarta from January 2011-December 2011 reached 59.2%, and premature rupture of membranes contributed 31.4% to sectio deliveries caesarea [11]. Research conducted by Sumelung in 2014 stated that premature rupture of membranes was one of the factors that increased the incidence of sectio caesarea at Liub Kendage Hospital in 2015 [12].

Based on the background above, the researchers wanted to research the relationship

between premature rupture of membranes and cesarean section delivery at UKI General Hospital in January-December 2019. The problem in this study was "Is there a relationship between premature rupture of membranes and cesarean delivery in UKI General Hospital period January-December 2019? The study aims to determine the incidence of premature rupture of membranes and sectio caesarea deliveries at UKI General Hospital from January to December 2019.

## **2. LITERATURE REVIEW**

The amniotic sac (amniotic sac), which lines the amniotic cavity, consists of the amnion and chorion, which are tightly bound. The amniotic membrane is a flexible but tough avascular tissue. These avascular structures have an important role in pregnancy in humans. In many obstetric cases, rupture of the membranes at a young gestational age is the most common cause of preterm birth. The inner membrane communicating with the amniotic fluid consists of a single layer of cuboidal epithelium derived from the embryonic ectoderm. This epithelium is tightly attached to the basement membrane, which is connected to the interstitial layer containing collagen I, III, and V. The outer part of the membrane is mesenchymal tissue derived from the mesoderm. This layer of the amnion is connected to the chorion leave. The amnion's inner lining is microvilli which transfer fluids and metabolism. This layer produces metalloproteinase-1 inhibitory substances [13].

Mesenchymal cells function to produce collagen so that the membrane becomes flexible and strong. In addition, these tissues produce the cytokines IL-6, IL-8, and MCP-1 (monocyte chemoattractant protein-1); This substance is useful against bacteria. In addition, the amniotic membranes produce vasoactive substances: endothelin-1 (a vasoconstrictor) and PHRP (parathyroid hormone-related protein), a vasorelaxant. Thus, the amniotic membrane regulates blood circulation and local vascular tone. The amniotic membrane also covers the umbilical cord. Some of the fluid will also come from diffusion in the umbilical cord. The rupture of the membranes is related to the strength of the membranes. In smokers and infections, the resistance of the membranes is weakening so that they break easily. In a normal pregnancy, there are few macrophages. At the time of birth, leukocytes will enter the amniotic fluid as a reaction to inflammation. There is no IL-1B in a

normal pregnancy, but IL-1B will be found in preterm labor.

It is related to the occurrence of the infection process. Since the beginning of pregnancy, amniotic fluid has been formed. Amniotic fluid is a protector and cushion for protection while supporting growth. Osmolarity, sodium, urea, and creatinine levels do not differ from the mother's serum levels, meaning that the levels in the amniotic fluid are the result of diffusion from the mother. Amniotic fluid contains many fetal cells (lanugo, vernix caseosa) [14]. Macroscopically, the amnion is a thin layer (only 20% of the thickness of the fetal membranes) and is a transparent (translucent) structure. This amniotic membrane can easily be separated from the underlying chorion because the amnion is never completely fused with the chorion and is only passively attached by internal pressure from the amniotic fluid [15]. The amnion lacks blood vessels and obtains its required nutrients and oxygen supplied directly by diffusion from the surrounding amniotic fluid, fetal surface vessels, or the decidua [16]. The thickness of the amnion varies greatly due to changes in the amount of mucin and fluid contained in the spongy lining, which ranges from 80 - 120  $\mu\text{m}$  [17].

The amniotic epithelium, the innermost amnion layer that leads to the fetus and surrounds the amniotic fluid, consists of a single layer of simple, squamous to cuboidal cells, with microvilli apical and attached to the basement membrane [18]. The amniotic epithelium secretes collagen types I, III, and IV, non-collagen glycoproteins such as laminin, nidogen, and fibronectin, forming a cell basement membrane thickness of nearly 1  $\mu\text{m}$ . The stromal compatum layer varies greatly in thickness, is tightly adherent, and cannot be separated from the basement membrane. This layer is a microfibril-dense cell-free homogeneous layer. Collagen types I, III, and V, and elastic fibers of the compact layer, are secreted by the mesenchymal cells in the fibroblast layer beneath the compact layer [19], which form the main burden-carrying fibrous framework of the amnion. The main tensile strength of the amnion is provided by collagen, which is arranged in one direction, i.e., parallel to the amniotic epithelium [20]. The compact layer and fibroblasts are difficult to differentiate histologically. The fibroblast layer beneath the compact layer is the most complex and thickest layer of the amnion, consisting of cellular tissue within a loose collagen network with non-collagenous glycoproteins, proteoglycans, and

type III collagen. Fibroblasts and macrophages (Hofbauer cells) have been identified within this layer. The spongy intermediate layer, also called the intermediate layer because it lies between the amnion and the chorion, is rich in proteoglycans and hyaluronate, which absorbs large amounts of water and forms a viscous layer that allows the amnion to slide over the chorion [21].

By this mechanism, this spongy coating makes it possible to absorb physical stress and to provide spontaneous short-term mechanical repair and defect closure systems. In addition, a circular fibrous structure is found in direct relation to the chorionic side, helping to adjust the lateral tension on the sliding surface between the amnion and chorion [22]. Occasionally, a potential space can be seen histologically in the spongy layer between the amnion and chorion. The space comprises loosely arranged collagen fibers that are sometimes oedematous and contain scattered fibroblasts and macrophages.

Development is between the embryonic cell mass and the trophoblast during the early implantation period. The small cells lining the inner surface of the trophoblast are called amniogenic cells, which are precursors of the amniotic epithelium. The amnion can be identified for the first time on embryo development's 7th or 8th day. It begins with the formation of a vesicle that develops into a small sac covering the dorsal surface of the embryo. As the amniotic sac enlarges, distention of the amniotic sac causes it to contact the interior surface of the chorion leave. Apposition of the chorion and amnion near the end of the first trimester. In young pregnancies, amniotic fluid is an ultrafiltrate of maternal plasma formed by the amnion cells. In the second trimester of pregnancy, amniotic fluid is formed by extracellular diffusion through the fetal skin, so its composition is similar to fetal plasma. Furthermore, after the second trimester, fetal skin horn substance formation occurs and blocks the diffusion of fetal plasma so that most of the amniotic fluid is formed by; amniotic cells and fetal urine. Fetal kidneys start excreting urine from the age of 12 weeks, and after reaching the age of 18 weeks, they can excrete urine as much as 7-14 cc/day. Term fetuses produce urine 27 cc/hour or 650 ccs daily [23]. Thus, the composition that forms amniotic fluid follows a postulate that the increase in amniotic fluid is not linear.

After the gestational age exceeds 12 weeks, what participates in forming amniotic fluid is; fetal kidneys (so that urea, creatinine, and uric acid are found), desquamation of fetal skin (skin cells, lanugo hair, vernix caseosa), secretions from the fetal lungs, transudate from the surface of the placental amnion, hormones or hormone-like substances in the amniotic fluid. Meanwhile, regulating amniotic fluid is very important to maintain the amount regularly. The following three important components influence the setting: production produced by the amnion cells, the amount of urine produced, and the amount of amniotic fluid swallowed by the fetus. Amniotic fluid regulation in term pregnancy includes the amount taken by the fetus  $\pm$  500-1000 ml, enters the lungs  $\pm$  170 ml, and from the umbilical cord and amnion  $\pm$  200-500 ml. At the same time, the amount of fluid released by the fetus into the amniotic cavity is from oral secretions  $\pm$  25 ml, secretions from the respiratory tract  $\pm$  170 ml, urine  $\pm$  800-1200 ml, and transmembrane from the amnion  $\pm$  10 ml. Thus, fetal urine appears to dominate the production of amniotic fluid, and the average regulation near term reaches 500 cc/day [24].

Amniotic fluid has an important role in supporting the process of pregnancy and childbirth. Throughout a normal pregnancy, the compartments of amniotic fluid provide space for the fetus to grow, move and develop. Without amniotic fluid, the uterus will contract and put pressure on the fetus; in cases where rupture of the membranes occurs early in the first trimester, the fetus may experience structural abnormalities, including facial distortion, reduction of limbs, and abdominal wall defects due to uterine compression [25]. Towards mid-pregnancy, the amniotic fluid becomes increasingly important for fetal development and growth, including lung development, so if there is insufficient amniotic fluid during mid-pregnancy, the fetus is often accompanied by pulmonary hypoplasia and continues to death. In addition, this fluid also has a protective role in the fetus. This liquid contains anti-bacterial agents and works to inhibit the growth of bacteria with potential pathogens. Amniotic fluid is a protective medium for the fetus to help with cervical dilation during labor and delivery. In addition, amniotic fluid also acts as a means of communication between the fetus and the mother. The maturity and readiness of the fetus for birth can be known from the fetal urine hormones excreted into the amniotic fluid [26].

Amniotic fluid can also be used as a diagnostic tool to see fetal growth and development abnormalities by conducting cell cultures or using a spectrometer. Other functions of amniotic fluid can also protect the fetus from trauma, as a medium for fetal musculoskeletal development, maintain fetal body temperature, equalize uterine pressure in parturition, clean the birth canal so that the baby is less infected, and maintain normal development and growth of the lungs and gastro tract intestinal [27]. Any wound on any body part will go through several stages: inflammation, proliferation, and maturation. Each phase involves many components and mechanisms, including the inflammatory, proliferative, and maturation phases [28].

Premature rupture of the membranes is a condition in which the membranes rupture before labor or in part. Premature rupture of membranes can occur in term or preterm pregnancies. Preterm Premature Rupture of the Membranes or premature rupture of membranes is the rupture of the membranes in patients with a gestational age of fewer than 37 weeks [29]. The incidence of premature rupture of membranes is 8% at term (> 37 weeks) and is associated with an increased risk of infection in the mother (chorioamnionitis, endometritis) and infection in the fetus (neonatal sepsis). Meanwhile, the incidence of preterm premature rupture of membranes is 3%, as a cause of 20-30% of preterm labor [30]. Two things broadly cause the causes of premature rupture of membranes, namely reduced membrane strength and increased intrauterine pressure or both factors. An infection can cause reduced membrane strength from the vagina and cervix. The factors influencing premature rupture of membranes are age, parity, smoking behavior, and history of premature rupture. Experiences experienced by mothers in labor with premature rupture of membranes can have a major influence on the mother when facing pregnancy conditions. The previous history of premature rupture of membranes is 2-4 times at risk of experiencing premature rupture of membranes again. In short, the pathogenesis of premature rupture of membranes results from a decrease in collagen content in the membrane, which triggers premature and preterm rupture of membranes. Women who have experienced premature rupture of membranes in pregnancies before delivery will be at greater risk in subsequent pregnancies than women who have never experienced premature rupture of membranes before because the composition of the

membranes decreases in subsequent pregnancies.

The rupture of membranes in labor is generally caused by repeated uterine contractions and stretching. The membranes rupture because, in certain areas, biochemical changes occur, which cause the inferior membranes to become brittle, not because the entire membranes are fragile. There is a balance between synthesis and degradation of the extracellular matrix. Changes in structure, cell number, and collagen catabolism cause collagen activity to change and cause the membranes to rupture. Collagen degradation is mediated by matrix metalloproteinases inhibited by tissue-specific and protease inhibitors. The membranes' integrity is due to a combination of low metalloproteinase activity and relatively higher concentrations of tissue inhibitors metalloproteinase-1. Microorganisms that infect the membranes of the membranes can form protease enzymes accompanied by an inflammatory response from the membranes, thereby affecting the balance of metalloproteinases and tissue inhibitors metalloproteinase-1, which causes a weakening of the tension of the membranes and rupture of the membranes hemolytic, and staphylococcus. In addition, closer to the delivery time, the balance between metalloproteinase and tissue inhibitors metalloproteinase-1 leads to proteolytic degradation of the extracellular matrix and fetal membranes. This proteolytic degradation activity increases towards delivery. The amniotic membranes are very strong in young pregnancies. In the third trimester, the membranes rupture easily. The weakening of the strength of the membranes is related to the enlargement of the uterus, uterine contractions, and fetal movements. In the last trimester, biochemical changes occur in the amniotic membranes so that the rupture of the membranes at term is physiological. Premature rupture of membranes in preterm pregnancy is caused by external factors such as infections that spread from the vagina [31].

Premature rupture of membranes greatly affects complications and length of labor. In the 1st stage of labor, the fetal membranes and the presenting part of the fetus act to open the upper part of the vagina. However, after the rupture of the membranes, the changes in the pelvic floor are all produced by the pressure exerted by the presenting part of the fetus so that the hydrostatic action of the fetal membranes causes

cervical dilatation. If the membranes have ruptured, the lowest part of the fetus that attaches to the cervix and forms the lower uterine segment functions the same. It will result in one of the complications of premature rupture of membranes, namely the long labor process, so sectio caesarea is the only option in this condition [32].

Another complication of mothers with premature rupture of membranes that causes morbidity and mortality is choriamnionitis. This infection causes inflammation of the amniotic fluid, amniotic membranes, or fetus, and this condition causes mothers with premature rupture of membranes to have symptoms of fever, fundal tenderness, tachycardia, fetal tachycardia, and smelly amniotic fluid. This condition is one of the complications feared to occur if a mother with premature rupture of membranes does not end her pregnancy immediately [33].

A study in Iran reported that out of 536 cases of premature rupture of membranes, 28% were performed by sectio caesarea. Other researchers concluded that sectio caesarea reached 26.6% in India in patients with premature rupture of membranes [34]. It is due to complications from premature membrane rupture, including fetal distress, malpresentation, narrow pelvis, failed induction, and the maternal complication of chorioamnionitis. Apart from that, other complications that led to the decision to perform sectio caesarea in mothers with premature rupture of membranes were complications of neonatal asphyxia and sepsis [35]. There is a relationship between fetal distress and decreased amniotic fluid volume. The more critical the fetus is. It was found that babies with premature rupture of membranes >12 hours had a 12 times greater risk of experiencing unfavorable outcomes, so it can be concluded that at term KPD, pregnancy termination should be carried out at least 6-12 hours from the onset of premature rupture of membranes.

### 3. RESEARCH METHODS

This type of research uses a descriptive analysis research method with a cross-sectional approach. This study used secondary data from UKI General Hospital medical records for the January-December 2019 period. The location for collecting research data was carried out in the UKI General Hospital medical records section. The time of execution, collection, and

implementation of the research was carried out from August 2020 – October 2020. The population for this study was patients who gave birth at the UKI General Hospital for the period January-December 2019. The sample was part of the population that had met the inclusion criteria, namely patients who gave birth on indications of Premature rupture of membranes at UKI General Hospital during January-December 2019. The Central Statistics Agency in Indonesia stated that the total population of Indonesia was 271,066,400 in 2019. The World Health Organization reported that as many as 17,655 mothers gave birth in Indonesia in 2019. The number of samples in this study was at least 65 people. Data collection was carried out at the UKI General Hospital medical record section. Furthermore, medical record data following the research variables will be recorded. All data obtained will be filtered again according to the inclusion and exclusion criteria set so that data is produced that will be used in the results of this study. Data collected through medical records are processed using the IBM SPSS (Statistical for Social Science) program for Windows edition 25.0 and the Microsoft Office Excel 2016 program.

### 4. RESULTS AND DISCUSSION

Data were obtained from the medical records of all patients with premature rupture of membranes delivered by cesarean section and normal delivery at UKI General Hospital for January-December 2019. The total sample was 65 patients.

Table 1 shows the characteristics of the study sample based on age, which found 15 patients (23.1%) aged <30 years, while 50 patients (76.9%) were aged ≥30 years.

**Table 1. Characteristics of the sample by age**

Age	Frequency	Percentage %
< 30 Tahun	15	23.1%
≥30 Tahun	50	76.9%
<b>Total</b>	<b>65</b>	<b>100%</b>

Table 2 shows the characteristics of the study sample based on parity, where there were 25 patients with primigravidas (38.5%), while multigravidas had 40 patients (61.5%).

Table 3 shows the characteristics of the study sample based on gestational age, which was found at gestational age <37 weeks in 12

patients (18.5%), while gestational age  $\geq 37$  weeks were found in 53 patients (81.5%).

**Table 2. Sample characteristics based on parity**

Parity	Frequency	Percentage %
Primigravida	25	38.5%
Multigravida	40	61.5%
<b>Total</b>	<b>65</b>	<b>100%</b>

**Table 3. Sample characteristics based on gestational age**

Gynecological age	Frequency	Percentage %
< 37 Weeks	12	18.5%
$\geq 37$ Weeks	53	81.5%
<b>Total</b>	<b>65</b>	<b>100%</b>

Table 4 shows the characteristics of the study sample based on BMI (Body Mass Index), where there were 17 patients (26.2%) with a BMI <30, while 48 patients (73.8%) had a BMI  $\geq 30$ .

Table 5 shows the characteristics of the study sample based on a history of premature rupture of membranes, in which 40 patients (61.5%) had a history of premature rupture of membranes, while 25 patients had never had premature rupture of membranes (38.5%).

**Table 4. Sample characteristics based on BMI (Body Mass Index)**

BMI	Frequency	Percentage %
< 30	17	26.2%
$\geq 30$	48	73.8%
<b>Total</b>	<b>65</b>	<b>100%</b>

**Table 5. Sample characteristics based on the history of premature rupture of membranes**

History of premature rupture of membranes	Frequency	Percentage %
Ever	40	61.5%
Never	25	38.5%
<b>Total</b>	<b>65</b>	<b>100%</b>

Table 6 shows the characteristics of the study sample based on the Apgar Score, where there were 46 patients (70.8%) with an APGAR SCORE <7, while 19 patients (29.2%) had an Apgar Score  $\geq 7$ .

Table 7 shows the characteristics of the study sample based on birth weight, in which 20 patients

(30.8%) had babies with birth weight <3 kg, while 45 patients (69.2%) had babies with birth weights  $\geq 3$  kg.

**Table 6. Characteristics of the sample based on the APGAR SCORE**

Apgar Score	Frequency	Percentage %
< 7	46	70.8%
$\geq 7$	19	29.2%
<b>Total</b>	<b>65</b>	<b>100%</b>

**Table 7. Sample characteristics based on birth weight**

Birth Weight	Frequency	Percentage %
< 3 Kg	20	30.8%
$\geq 3$ Kg	45	69.2%
<b>Total</b>	<b>65</b>	<b>100%</b>

Table 8 shows the characteristics of the study sample based on the gestational age of mothers who experienced premature rupture of membranes and the type of delivery performed. It was found that mothers with gestational age <37 weeks experienced premature rupture of membranes and performed sectio caesarea in 7 patients (10.76%), whereas mothers with gestational age  $\geq 37$  weeks experienced premature rupture of membranes but did not perform a cesarean section in 5 patients (7.69%). Meanwhile, 50 patients (76.92%) had premature rupture of membranes, and 50 patients (76.92%) had premature rupture of membranes and 37 weeks of gestational age  $\geq 37$  weeks of mothers who had premature rupture of membranes without having a caesarean section. (4.63%), and the P-value is 0.001.

The study results found that the age group that experienced premature rupture of membranes was the most in the age group  $\geq 30$  years, with as many as 50 patients (76.9%). It follows the theory where according to Sudarto, a mother who gives birth at the age of 35 and over can affect her reproductive system because her reproductive organs have begun to reduce their ability and elasticity in accepting pregnancy, affecting the process of embryogenesis. The amniotic membranes will be thinner, facilitating premature rupture [36]. Based on the results of the study, it was found that a woman with multigravida experienced premature rupture of membranes in 40 patients (61.5%). It is following the theory according to Kacerovsky, women who have given birth several times accompanied by

close pregnancy intervals can be at risk of experiencing premature rupture of membranes in their next pregnancy. It is because pregnancies are too frequent, affecting the embryogenesis process. The amniotic membranes become thinner, so they break easily prematurely. In addition, there is also a theory that the more parity, the easier it is for amnion infection to occur due to damage to the cervical structure in previous deliveries [37].

Based on the study's results, 53 pregnant women experienced premature rupture of membranes when their gestational age was  $\geq 37$  weeks (81.5%). According to Popowski, it follows the theory that most premature rupture of membranes occurs at gestational age  $\geq 37$  weeks. Premature rupture of membranes occurs nearing delivery due to an increase in matrix metalloproteinase which tends to cause the membranes to rupture, in addition to an increase in matrix metalloproteinase, uterine enlargement, uterine contractions, and fetal movements occur [38].

From the results of the study, it was found that pregnant women who had a BMI (Body Mass Index)  $\geq 30$  had premature rupture of membranes, and 48 patients (73.8%) had a cesarean section. An increase in Body Mass Index  $\geq 30$  has a great risk for a normal delivery because, in a normal delivery, there will be a delay in the first stage of the active phase and the second stage. After all, the mother who is obese will have a narrow birth canal due to the accumulation of fat, making it difficult for the baby to come out, and a sectio caesarea must be carried out immediately [39].

The history that mothers have experienced in labor with the incidence of premature rupture of membranes can have a big influence on the mother when facing pregnancy conditions. Based on the amount of research data, it was found that 40 patients (61.5%) had a history of premature

rupture of membranes in previous deliveries. According to Morgan, it follows the theory that patients with a previous history of premature rupture of membranes are at risk of 2-4 times experiencing premature rupture of membranes again. In short, the pathogenesis of premature membrane rupture results from a decrease in collagen content in the membrane, which triggers premature rupture membranes.

Women who have experienced premature rupture of membranes during pregnancy or before delivery are at greater risk in their next pregnancy than women who have never experienced premature rupture of membranes before because the composition of the membranes becomes fragile. The collagen content decreases in subsequent pregnancies [40]. The APGAR SCORE is a practical method systematically used to assess a baby as soon as possible after birth. What is assessed is heart rate, respiratory effort, muscle tone, skin color, and reaction to stimulation (response to stimuli), namely by inserting a catheter into the nostril after the airway has been cleaned when the new baby is born.

From the results of this assessment, it can be seen whether the baby is normal (APGAR score 7-10), moderate-mild asphyxia (APGAR score 4-6), or the baby has severe asphyxia (APGAR score 0-3). The characteristics of the sample based on the APGAR SCORE showed that patients who gave birth with premature rupture of membranes and their babies had an APGAR SCORE  $<7$ . There were 46 patients (70.8%). It follows the theory according to Ni Wayan and Melania, where with the rupture of the membranes, there will be prolapse of the funiculi / descent of the umbilical cord, which will cause the neonatorum to experience hypoxia and asphyxia (lack of oxygen in the baby) which will result in compression of the umbilical cord, respiratory distress. The baby will be at risk of having a low APGAR SCORE ( $<7$ ) [41].

**Table 8. The relationship between premature rupture of membranes and sectio caesarea delivery**

Gynecological Age	Type of childbirth				Total	p-value
	Section	n (%)	No section	n (%)		
Premature rupture of membranes	< 37 Weeks	7	10,76%	5	7,69%	12(18.45%)
	$\geq 37$ Weeks	50	76,92%	3	4,63%	
<b>Total</b>		57	87,68%	8	12,32%	65 (100%)



Mechanical stretching, such as in a large baby's weight, can cause stretching of the membranes and can cause premature rupture of the membranes. It is also supported by the data results where there is an increase in the number of patients who experience premature rupture of membranes with a baby weighing > 3.5 kg by 1.6%, while babies weighing > 4 Kg were 2.4%. Based on the study's results, with birth weight  $\geq$  3 Kg and experiencing premature rupture of membranes were 45 patients (69.2%). It follows the theory according to Zelop, where mechanically stretching the fetal membranes will increase the production of prostaglandin E2 and Interleukin-8 in the amnion, also increasing the activity of MMP-1 in the membranes. Interleukin-8 produced from amnion and chorion cells is chemotactic towards neutrophils and stimulates collagenase activity, which disrupts the balance of synthesis and degradation of the extracellular matrix, which ultimately causes the rupture of the membranes.

Prostaglandin E2 increases uterine irritability, decreases membrane collagen synthesis, and increases the production of MMP-1 and MMP-3 by fibroblasts. The production of interleukin-8 and prostaglandin E2 from the amnion indicates that biochemical changes in the membranes can be initiated by physical forces or by stretching the membranes, suggesting that mechanical forces cause premature rupture of membranes [42].

The results of the research on the relationship between premature rupture of membranes and sectio caesarea delivery obtained a value of  $P = 0.001$  ( $P < 0.05$ ) which means there is a relationship between premature rupture of membranes and sectio caesarea delivery where this is following the theory, according to Barbara, pregnant women with indications of ruptured membranes Early pregnancy will cause the amount of amniotic fluid to decrease and the risk of having a normal delivery. Babies born will also have an increased risk of morbidity and mortality. The risk of infection in the mother is also quite high, so it will increase morbidity and mortality, so the action of cesarean section is expected to reduce maternal morbidity and mortality and improve infant outcomes.

## 5. CONCLUSION

From the results of the research that was described in the previous chapter, the following conclusions were drawn: a) the prevalence of premature rupture of membranes at UKI General

Hospital carrying out sectio caesarea in the January-December 2019 period was 57 patients (87.68%), while those who did not perform sectio caesarea in 8 patients (12.32%), and b) based on the Chi-square test, there was a significant relationship between premature rupture of membranes and sectio caesarea ( $P=0.001$ ). Thus, it is hoped that health workers and health institutions as health workers can help educate patients about the factors that can cause premature rupture of membranes so that patients can take preventive action to prevent premature rupture membranes. In addition, it is recommended that patients avoid factors that can cause premature rupture of a membrane that doctors or health workers have told, and the patient's family can support patients to remind them to avoid factors that can cause premature rupture of membranes.

## CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

## ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Caughey AB, Robinson JN, Norwitz ER. Contemporary diagnosis and management of preterm premature rupture of membranes. *Reviews in Obstetrics and Gynecology*. 2008;1(1):11.
2. Menon R, Richardson LS. Preterm prelabor rupture of the membranes: A disease of the fetal membranes. In *Seminars in Perinatology*. 2017;41(7): 409-419. WB Saunders.
3. Fitriyani F, Yuliana Lubis Y. Determinant factors in premature rupture of membranes. *Health Media Journal*. 2018; 11(1):053-61.
4. Syarwani TI, Tendean HM, Wantania JJ. Overview of Premature Ruptured Membranes (PROM) in Prof. Dr. RD

- Kandou Manado in 2018. Medical Scope Journal. 2020;1(2).
5. Susan Blossia E. A Prospective Cross Sectional study on CRP levels in antenatal women with prelabor rupture of membrane and crp levels in women with normal labor (Doctoral dissertation, Christian Medical College, Vellore).
  6. Rohmawati N, Wijayanti Y. Premature rupture of membranes at Ungaran Regional General Hospital. HIGEIA (Journal of Public Health Research and Development). 2018;2(1):23-32.
  7. Venancyana MC, Indrayani T, Lubis R. Analysis of factors affecting pregnant mothers in making decisions for caesarean section childbirth plan at Tc Hillers Hospital, Sikka Regency, Year 2022. International Journal of Multi Science. 2022;3(02):7-17.
  8. Waang WW, Kusumawati PD. Analysis of Deep Breath Relaxation Techniques for Post SC Patients with Pain. Journal of Health Science Community. 2022;3(2):97-101.
  9. BP Statistics. Indonesia Demographic and Health Survey Report.
  10. Suparmi S, Kusumawardhani N, Susiloretni KA. Sub-national inequality of caesarean sections in urban-rural areas of Indonesia. In Proceedings of the International Conference on Public Health. 2018;4(1):23-32.
  11. Hardiyanti R. Use of prophylactic antibiotics in section caesarea patients. Journal of Health Science and Physiotherapy. 2020;2(1):96-105.
  12. Ulfa EN. Factors related to delivery by sectio caesarea at the Karsa Husada Batu general hospital in 2020 (Doctoral dissertation, Maulana Malik Ibrahim State Islamic University).
  13. [Machtejeviené E. Peptide pattern of amniotic fluid and its correlation with protein composition of fetal membranes: The search for new potential biomarkers to predict preterm premature rupture of membranes: [Rankraštis]: doctoral dissertation: Biomedical Sciences Medicine (06B).
  14. Brace RA, Anderson DF, Cheung CY. Regulation of amniotic fluid volume: mathematical model based on intramembranous transport mechanisms. American Journal of Physiology-Regulatory, Integrative and Comparative Physiology. 2014;307(10):R1260-73.
  15. Cohen MC, Scheimberg I, Hutchinson JC. Anatomy and pathology of the placental membranes. Benirschke's Pathology of the Human Placenta. 2022;281-343.
  16. Founds SA, Baker B. Fetal and Placental Development and Functioning. Core Curriculum for Maternal-Newborn Nursing E-Book. 2022;28.
  17. POLZIN WJ, BRADY K. Mechanical factors in the etiology of premature rupture of the membranes. Clinical Obstetrics and Gynecology. 1991;34(4):702-14.
  18. Mamede AC, Carvalho MJ, Abrantes AM, Laranjo M, Maia CJ, Botelho MF. Amniotic membrane: From structure and functions to clinical applications. Cell and Tissue Research. 2012;349:447-58.
  19. Country IK, KFM SO. Matrix Metalloproteinases in premature ruptured membrane. Deepublish; 2021.
  20. Yunus M, Sos S, Hatijar S. Textbook of Midwifery Care in Pregnancy.
  21. Macedo BM. Exploração das Membranas Fetais para Aplicações de Engenharia dos Tecidos (Doctoral dissertation).
  22. Haller C. Evaluation of human fetal membrane sealing (Doctoral dissertation, University of Zurich).
  23. Ferselina L. Case study of handling newborns with cloudy amniotic water in Mrs. Vdi Pmb Komariah, Sst Lampung Selatan (Doctoral dissertation, Tanjungkarang Health Polytechnic).
  24. Fahimah WF. Literature Review: Factors Causing Premature Rupture of Water.
  25. Trance MG. Continuing Midwifery Care for Mrs. EB at the Kaputu Community Health Center April 13 to June 10 2019 (Doctoral dissertation, Kupang Ministry of Health Polytechnic).
  26. Liley AW. Disorders of amniotic fluid. In Fetal-placental disorders 1972;157-206. Academic Press.
  27. Henny Sulistyawati SS, Zulfa Hanum SS, Keb M, Meikawati PR, SiTS S, Rosdianto NO, ST S, Keb M, Muayah SS, Ety Aprianti SK. Textbook for Postpartum DIII Midwifery Volume I. Mahakarya Citra Utama Group; 2022.
  28. Monika P, Waiker PV, Chandraprabha MN, Rangarajan A, Murthy KN. Myofibroblast progeny in wound biology and wound healing studies. Wound Repair and Regeneration. 2021;29(4):531-47.
  29. Ma'roef M, Putri AP. The relationship of leukocyturia on the incidence of prominent rupture of the membrane in preterm

- Pregnancy at The General Hospital of The University of Muhammadiyah Malang. Medical Science. 2019;15(2):139-45.
30. Sari DS. Relationship between multiple pregnancy and fetal abnormalities with premature rupture of the membranes at the Prabumulih Regional General Hospital in 2019. *Abdurahman Health Journal*. 2020; 9(2):56-63.
  31. Margono RS, Sukrisno A, Nugrohowati N, Lestari W. Relationship between a premature rupture of membranes and the increase of leucocyte levels in pregnant women during the COVID-19 pandemic. *Journal of Traditional Midwifery and Health*. 2021:127-34.
  32. Putri RS. Application of the Pelvic Rocking Technique on the Gymball Against the Progress of 1st Stage Labor Ny. T At PMB Eliana Putriani, Str. South Lampung Regency (Doctoral dissertation, Tanjungkarang Health Polytechnic).
  33. Sprong KE, Mabenge M, Wright CA, Govender S. Ureaplasma species and preterm birth: Current perspectives. *Critical Reviews in Microbiology*. 2020;46(2):169-81.
  34. Ulfa EN. Factors related to delivery by sectio caesarea at the Karsa Husada Batu general hospital in 2020 (Doctoral dissertation, Maulana Malik Ibrahim State Islamic University).
  35. Marlyana DM, Nurwahyuni A, Wibowo A. Neonate outcome and response time for emergency cesarean section: A scoping review. *Journal of Medicine and Health: Scientific Publication of the Faculty of Medicine, Sriwijaya University*. 2023; 10(1):125-40.
  36. Motiec TD, Trisnawati I, Handayani IF, Darwanti J. Description of the case of Premature Ruptured Membranes in Mothers at the Purwasari Karawang Maternity Hospital in 2020 (Doctoral dissertation, Health Polytechnic of the Ministry of Health, Bandung).
  37. Wahyuni R. Factors causing premature rupture of membranes in women giving birth at the general hospital in the Sumbawa Besar area. *Journal of Health and Science*. 2020;3(2):26-33.
  38. Dea D, Riana E, Khairunisa D. Midwifery care for women giving birth with premature rupture of membranes in the independent practice of Midwife Ida Apianti Pontianak (Doctoral dissertation, Polytechnic'Aisyiyah Pontianak).
  39. Mendez-Figueroa H, Bicocca MJ, Bhalwal AB, Wagner SM, Chauhan SP, Bartal MF. Preterm cesarean delivery for nonreassuring fetal heart rate tracing: Risk factors and predictability of adverse outcomes. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2022;276:207-12.
  40. IM's daughter, Ismiyatun N. Early detection of high-risk pregnancies. *JKM (Journal of Public Health) Main Scholar*. 2020;8(1):40-51.
  41. Yuliani NW, Wahyuningsih M. Differences in apgar values in the first minutes of newborns with premature rupture of membranes and normal rupture of Membranes in Yogyakarta City Hospital. In *Proceedings of the National Seminar on Multidisciplinary Sciences*. 2019;1(1).
  42. Leimert KB, Xu W, Princ MM, Chemtob S, Olson DM. Inflammatory amplification: A central tenet of uterine transitions for labor. *Frontiers in Cellular and Infection Microbiology*. 2021;11: 660983.

© 2023 Sihombing et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/97587>