



C_n -E- super Magic Graceful Labeling of Some Special Graphs

M. Sindhu ^{a++*} and S. Chandra Kumar ^b

^aDepartment of Mathematics, Excel Engineering College (Autonomous), Komarapalayam-637303, Namakkal, Tamil Nadu, India.

^bDepartment of Mathematics, Scott Christian College (Autonomous), Nagercoil-629003, Tamil Nadu, India.

Authors' contributions

This work was carried out in collaboration between boht authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJOM/2023/v19i7677

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98052>

Received: 25/01/2023

Accepted: 28/03/2023

Published: 21/04/2023

Original Research Article

Abstract

A graph G possess an H -covering when each edge in $E(G)$ pertaining to a subgraph of G isomorphic to H . This graph G is H -magic if there exists a total labeling $f : V(G) \cup E(G) \rightarrow \{1, 2, \dots, p + q\}$ such that for each subgraph H' of G isomorphic to H , $\sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e) = M$ is a constant. An H -E-super magic graceful labeling (H -E-SMGL) is a bijective function $f : V(G) \cup E(G) \rightarrow \{1, 2, \dots, p + q\}$ with $f(E(G)) = \{1, 2, \dots, q\}$ so that $\sum_{v \in V(H')} f(v) - \sum_{e \in E(H')} f(e) = M$ for few positive integer M . Herein, we examine the C_n -E-SMGL of some graphs.

Keywords: H -covering; H -magic labeling; H -E-super magic labelling; H -E- super magic graceful labeling.

⁺⁺AssistantProfessor

Correspondingauthor : E – mail : msindhu0387@gmail.com;

Asian Res. J. Math., vol. 19, no. 7, pp. 38 – 46, 2023

AMS subject classification code: 05C78.

1 Introduction

All graphs considered in this article are finite, simple and undirected. The vertex set and edge set of a graph G is represented as $V(G)$ and $E(G)$ correspondingly, $p = |V|$ and $q = |E|$. A graph labeling is a map that takes graph elements to numbers (typically integers). Various classes of labelings has been introduced by several experts. An excellent analysis of graph labelings is glimpsed in [1].

During 1963, Sedláček [2] described magic labeling in graphs. A graph G is magic when the edges of G usually labeled with $\{1, 2, \dots, q\}$ such that the sum over the labels of all edges incident with any vertex is equal [3]

$$\sum_{v \in N(v)} f(uv) = M.$$

A covering of G is a family of subgraphs H_1, H_2, \dots, H_h so that each edge of $E(G)$ pertaining to at least one of the subgraphs $H_i, 1 \leq i \leq h$. This results that G possess an (H_1, H_2, \dots, H_h) covering. When each H_i is isomorphic to the graph H , then G have an H -covering. Assume that G have an H -covering. A total labeling is a bijective function f from $V(G) \cup E(G)$ to $\{1, 2, 3, \dots, |V(G)| + |E(G)|\}$ is named an H -magic labeling of G if there exists a positive integer M (termed the magic constant) so that for every subgraph H' of G isomorphic to H , $\sum_{v \in V(H')} f(v) + \sum_{e \in E(H')} f(e) = M$. A graph which possess such a labeling is termed H -magic. The function f is named as H - E -super magic labeling when $f(E(G)) = \{1, 2, \dots, q\}$.

The concept of H -magic labeling was explained by Gutierrez and Llado [4].

Llado and Moragas [5] explored few C_n -supermagic graphs.

Rosa [6] initiated a labeling known as β -valuation. Golomb [7] named that labeling as graceful. An one to one function f from the vertices of G to $\{0, 1, 2, \dots, q\}$ is named as graceful labeling of G when every edge uv is labeled as $|f(u) - f(v)|$, the resultant edge labels are different.

To acquire more knowledge regarding H - E -super magic graphs, read [8].

In 2019, Sindhu Murugan and S. Chandra Kumar [9] initiated an H - E -super magic graceful labeling (H - E -SMGL). An H - E -SMGL is a bijective function f from $V(G) \cup E(G)$ to $\{1, 2, \dots, p + q\}$ with $f(E(G)) = \{1, 2, \dots, q\}$ and $\sum_{v \in V(H')} f(v) - \sum_{e \in E(H')} f(e) = M$ for few positive integer M . Herein, we examine C_n - E -SMGL of some families of graphs.

There are so many types of magic labelings in graphs, defined and studied by various authors [10, 11, 12, 13, 14, 15, 16, 17]

2 C_n - E -Super Magic Graceful Graphs

Theorem 2.1. *Let $n \geq 5$ be an odd integer. Then the wheel graph W_n is C_3 - E -SMGL with magic constant $\frac{9n+5}{2}$.*

Proof. Denote the vertices of n -cycle of the wheel W_n as a_1, a_2, \dots, a_n and its central vertex by r . We define a total labeling $f : V(W_n) \cup E(W_n) \rightarrow \{1, 2, 3, \dots, 3n + 1\}$ as follows:

$$f(v) = \begin{cases} 2n + 1 & \text{if } v = r \\ 2n + 2 & \text{if } v = a_1 \\ 2n + \frac{i+3}{2} & \text{if } v = a_i, i \text{ is odd for } 3 \leq i \leq n \\ \frac{5n+3+i}{2} & \text{if } v = a_i, i \text{ is even for } 2 \leq i \leq n - 1 \end{cases}$$

and

$$f(e) = \begin{cases} i & \text{if } e = ra_i \text{ for } 1 \leq i \leq n \\ 2n + 1 - i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq n - 1 \\ n + 1 & \text{if } e = a_n a_1. \end{cases}$$

Now, we prove that f is a $C_3 - E$ -SMGL of W_n .

Let C_3^i for $1 \leq i \leq n$ be the subcycle of W_n with $V(C_3^i) = \{a_i : 1 \leq i \leq n\} \cup \{r\}$ and $E(C_3^i) = \{a_i a_{i \oplus n 1} : 1 \leq i \leq n\} \cup \{ra_i : 1 \leq i \leq n\} \cup \{ra_{i \oplus n 1} : 1 \leq i \leq n\}$.

Case 1: Suppose $i = 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_1) + f(a_2) - [f(a_1 a_2) + f(ra_1) + f(ra_2)] \\ &= [2n + 1] + [2n + 2] + [\frac{5n+5}{2}] - [2n + 1 + 2] = \frac{9n+5}{2}. \end{aligned}$$

Case 2: Suppose i is even for $2 \leq i \leq n - 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_i) + f(a_{i+1}) - [f(a_i a_{i+1}) + f(ra_i) + f(ra_{i+1})] \\ &= [2n + 1] + [\frac{5n+3+i}{2}] + [2n + 2 + \frac{i}{2}] - [2n + 1 - i + i + i + 1] = \frac{9n+5}{2}. \end{aligned}$$

Case 3: Suppose i is odd for $3 \leq i \leq n - 2$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_i) + f(a_{i+1}) - [f(a_i a_{i+1}) + f(ra_i) + f(ra_{i+1})] \\ &= [2n + 1] + [2n + \frac{i+3}{2}] + [\frac{5n+4+i}{2}] - [2n + 1 - i + i + i + 1] = \frac{9n+5}{2} \end{aligned}$$

Case 4: Suppose $i = n$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_n) + f(a_1) - [f(a_n a_1) + f(ra_1) + f(ra_n)] \\ &= [2n + 1] + [\frac{5n+3}{2}] + [2n + 2] - [n + 1 + 1 + n] = \frac{9n+5}{2}. \end{aligned}$$

The graph W_n is $C_3 - E$ -SMG with magic constant $\frac{9n+5}{2}$. □

Example 2.2. The Wheel W_7 admits C_3 -E-SMGL with magic constant 34.

Denote the vertices of n -cycle of the wheel W_n as a_1, a_2, \dots, a_7 and its central vertex by r . Define $f : V(W_7) \cup E(W_7) \rightarrow \{1, 2, 3, \dots, 22\}$ as follows:

$$f(v) = \begin{cases} 15 & \text{if } v = r \\ 16 & \text{if } v = a_1 \\ 14 + \frac{i+3}{2} & \text{if } v = a_i, i \text{ is odd for } 3 \leq i \leq 7 \\ 19 + \frac{i}{2} & \text{if } v = a_i, i \text{ is even for } 2 \leq i \leq 6 \end{cases}$$

and

$$f(e) = \begin{cases} i & \text{if } e = ra_i \text{ for } 1 \leq i \leq 7 \\ 15 - i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq 6 \\ 8 & \text{if } e = a_7 a_1. \end{cases}$$




Fig. 1. C_3 -E-SMGL of W_7

To prove that f is a $C_3 - E$ -SML of W_7 .

Let C_3^i for $1 \leq i \leq n$ be the subcycle of W_n with $V(C_3^i) = \{a_i : 1 \leq i \leq 7\} \cup \{r\}$ and $E(C_3^i) = \{a_i a_{i \oplus n} : 1 \leq i \leq 7\} \cup \{ra_i : 1 \leq i \leq 7\} \cup \{ra_{i \oplus 7} : 1 \leq i \leq 7\}$.

Case 1: Suppose $i = 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^1)} f(v) - \sum_{e \in E(C_3^1)} f(e) = f(r) + f(a_1) + f(a_2) - [f(a_1 a_2) + f(ra_1) + f(ra_2)] \\ &= [15] + [16] + [20] - [14 + 1 + 2] = 34. \end{aligned}$$

Case 2: Suppose i is even for $2 \leq i \leq 6$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_i) + f(a_{i+1}) - [f(a_i a_{i+1}) + f(ra_i) + f(ra_{i+1})] \\ &= [15] + [19 + \frac{i}{2}] - [16 + \frac{i}{2}] - [15 - i + i + i + 1] = 34. \end{aligned}$$

Case 3: Suppose i is odd for $3 \leq i \leq 5$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_i) + f(a_{i+1}) - [f(a_i a_{i+1}) + f(ra_i) + f(ra_{i+1})] \\ &= [15] + [14 + \frac{i+3}{2}] + [\frac{39+i}{2}] - [15 - i + i + i + 1] = 34 \end{aligned}$$

Case 4: Suppose $i = 7$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(r) + f(a_n) + f(a_1) - [f(a_n a_1) + f(r a_1) + f(r a_n)] \\ &= [15] + [19] + [16] - [8 + 1 + 7] = 34. \end{aligned}$$

The graph W_n is $C_3 - E$ -SMG with magic graceful constant 34.

Theorem 2.3. *Let $n \geq 1$ be an integer. Then the Ladder graph $L_n = P_2 \times P_n$ admits C_4 -E-SMGL with magic constant $9n + 4$.*

Proof. Let $V(L_n) = \{a_i, b_i : 1 \leq i \leq n\}$ and $E(L_n) = \{a_i a_{i+1}, b_i b_{i+1} : 1 \leq i \leq n-1\} \cup \{a_i b_i : 1 \leq i \leq n\}$ be the vertex set and the edge set of L_n respectively.

We define a total labeling $f : V(L_n) \cup E(L_n) \rightarrow \{1, 2, \dots, 5n - 2\}$ as follows:

$$f(v) = \begin{cases} 2n + i + 3 & \text{if } v = a_i \text{ for } 1 \leq i \leq n \\ 5n - i - 1 & \text{if } v = b_i \text{ for } 1 \leq i \leq n \end{cases}$$

$$f(e) = \begin{cases} i & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq n \\ 2n - i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq n - 1 \\ 3n - i - 1 & \text{if } e = b_i b_{i+1} \text{ for } 1 \leq i \leq n - 1. \end{cases}$$

Now, we prove that f is a $C_4 - E$ -SMGL of L_n .

Let C_4^i for $1 \leq i \leq n - 1$ be the subcycle of L_n with $V(C_4^i) = \{a_i, b_i : 1 \leq i \leq n\}$ and $E(C_4^i) = \{a_i a_{i+1} : 1 \leq i \leq n - 1\} \cup \{b_i b_{i+1} : 1 \leq i \leq n - 1\} \cup \{a_i b_i : 1 \leq i \leq n\}$.

Suppose $1 \leq i \leq n - 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_4^i)} f(v) - \sum_{e \in E(C_4^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_i) + f(b_{i+1}) - [f(a_i b_i) + f(a_{i+1} b_{i+1}) + f(a_i a_{i+1}) + \\ & f(b_i b_{i+1})] \\ &= [2n + i + 3] + [2n + i + 4] + [5n - i - 1] + [5n - i - 2] - [i + i + 1 + 2n - i + 3n - i - 1] = 9n + 4. \end{aligned}$$

The graph L_n is $C_4 - E$ -SMG with magic constant $9n + 4$. \square

Example 2.4. *The Ladder graph $L_5 = P_2 \times P_5$ admits C_4 -E-SMGL with magic constant 49.*

Let $V(L_5) = \{a_i, b_i : 1 \leq i \leq 5\}$ and $E(L_5) = \{a_i a_{i+1}, b_i b_{i+1} : 1 \leq i \leq 4\} \cup \{a_i b_i : 1 \leq i \leq 5\}$ be the vertex set and the edge set of L_5 respectively.

Define $f : V(L_5) \cup E(L_5) \rightarrow \{1, 2, \dots, 23\}$ as follows:

$$f(v) = \begin{cases} 13 + i & \text{if } v = a_i \text{ for } 1 \leq i \leq 5 \\ 24 - i & \text{if } v = b_i \text{ for } 1 \leq i \leq 5 \end{cases}$$

and

$$f(e) = \begin{cases} i & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq 5 \\ 10 - i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq 4 \\ 14 - i & \text{if } e = b_i b_{i+1} \text{ for } 1 \leq i \leq 4. \end{cases}$$




Fig. 2. C_4 -E-SMGL of L_5

To prove that f is a $C_4 - E$ -SMGL of L_5 .

Let C_4^i for $1 \leq i \leq 4$ be the subcycle of L_5 with $V(C_4^i) = \{a_i, b_i : 1 \leq i \leq 5\}$ and $E(C_4^i) = \{a_i a_{i+1} : 1 \leq i \leq 4\} \cup \{b_i b_{i+1} : 1 \leq i \leq 4\} \cup \{a_i b_i : 1 \leq i \leq 5\}$.

Suppose $1 \leq i \leq 4$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_4^i)} f(v) - \sum_{e \in E(C_4^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_i) + f(b_{i+1}) - [f(a_i b_i) + f(a_{i+1} b_{i+1}) + f(a_i a_{i+1}) + f(b_i b_{i+1})] \\ &= [13 + i] + [14 + i] + [24 - i] + [23 - i] - [i + i + 1 + 10 - i + 14 - i] = 49. \end{aligned}$$

Thus the graph L_5 is $C_4 - E$ -SMG with magic constant 49.

Theorem 2.5. Let $n \geq 2$ be an integer. Then the triangular Ladder TL_n admits C_3 -E-SMGL with magic constant $M = 10n - 5$.

Proof. Let $V(TL_n) = \{a_i, b_i : 1 \leq i \leq n\}$ and $E(TL_n) = \{a_i a_{i+1}, b_i b_{i+1} : 1 \leq i \leq n - 1\} \cup \{a_i b_i : 1 \leq i \leq n\} \cup \{a_i b_{i+1} : 1 \leq i \leq n - 1\}$ be the vertex set and the edge set of TL_n respectively.. We define a total labeling $f : V(TL_n) \cup E(TL_n) \rightarrow \{1, 2, \dots, 6n - 3\}$ as follows:

$$f(v) = \begin{cases} 4n + 2i - 3 & \text{if } v = a_i \text{ for } 1 \leq i \leq n \\ 4n + 2i - 4 & \text{if } v = b_i \text{ for } 1 \leq i \leq n \end{cases}$$

and

$$f(e) = \begin{cases} 2i - 1 & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq n \\ 2n + 2i - 2 & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq n - 1 \\ 2n + 2i - 1 & \text{if } e = b_i b_{i+1} \text{ for } 1 \leq i \leq n - 1 \\ 2i & \text{if } e = a_i b_{i+1} \text{ for } 1 \leq i \leq n - 1. \end{cases}$$

To prove that f is a $C_3 - E$ -SMGL of TL_n .

Let C_3^i for $1 \leq i \leq n - 1$ be the subcycle of TL_n with $V(C_3^i) = \{a_i : 1 \leq i \leq n\} \cup \{b_i : 1 \leq i \leq n\}$ and $E(C_3^i) = \{a_i a_{i+1} : 1 \leq i \leq n - 1\} \cup \{b_i b_{i+1} : 1 \leq i \leq n - 1\} \cup \{a_i, b_i : 1 \leq i \leq n\}$. Suppose $1 \leq i \leq n - 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_{i+1}) - [f(a_i a_{i+1}) + f(a_{i+1} b_{i+1}) + f(b_{i+1} a_i)] = \\ &= [4n + 2i - 3] + [4n + 2i - 1] + [4n + 2i - 2] - [2n + 2i - 2 + 2i + 1 + 2i] = 10n - 5. \end{aligned}$$

The graph TL_n is $C_3 - E$ -SMG with magic constant $10n - 5$. □

Example 2.6. The triangular Ladder TL_5 admits C_3 -E-SMGL with magic constant $M = 45$.

Let $V(TL_5) = \{a_i, b_i : 1 \leq i \leq 5\}$ and $E(TL_5) = \{a_i a_{i+1}, b_i b_{i+1} : 1 \leq i \leq 4\} \cup \{a_i b_i : 1 \leq i \leq 5\} \cup \{a_i b_{i+1} : 1 \leq i \leq 4\}$ be the vertex set and the edge set of TL_5 respectively.

Define $f : V(TL_5) \cup E(TL_5) \rightarrow \{1, 2, \dots, 27\}$ as follows:

$$f(v) = \begin{cases} 17 + 2i & \text{if } v = a_i \text{ for } 1 \leq i \leq 5 \\ 16 + 2i & \text{if } v = b_i \text{ for } 1 \leq i \leq 5 \end{cases}$$

and

$$f(e) = \begin{cases} 2i - 1 & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq 5 \\ 8 + 2i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq 4 \\ 9 + 2i & \text{if } e = b_i b_{i+1} \text{ for } 1 \leq i \leq 4 \\ 2i & \text{if } e = a_i b_{i+1} \text{ for } 1 \leq i \leq 4. \end{cases}$$




Fig. 3. C_3 -E-SMGL of triangular ladder TL_5

To prove that f is a $C_3 - E$ -SMGL of TL_5 .

Let C_3^i for $1 \leq i \leq 4$ be the subcycle of TL_5 with $V(C_3^i) = \{a_i : 1 \leq i \leq 5\} \cup \{b_i : 1 \leq i \leq 5\}$ and $E(C_3^i) = \{a_i a_{i+1} : 1 \leq i \leq 4\} \cup \{b_i b_{i+1} : 1 \leq i \leq 4\} \cup \{a_i, b_i : 1 \leq i \leq 5\}$. Suppose $1 \leq i \leq n - 1$.

$$\begin{aligned} \text{Then } M &= \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_{i+1}) - [f(a_i a_{i+1}) + f(a_{i+1} b_{i+1}) + f(b_{i+1} a_i)] \\ &= [17 + 2i] + [17 + 2i + 2] + [16 + 2i + 2] - [2i + 1 + 8 + 2i + 2i] = 45. \end{aligned}$$

Thus the graph TL_5 is $C_3 - E$ -SMG with magic constant 45.

Theorem 2.7. Let $n \geq 2$ be an integer. Then the triangular snake graph Δ_n admit C_3 -E-SMGL with magic constant $M = 7n + 2$.

Proof. Let $V(\Delta_n) = \{a_i : 1 \leq i \leq n + 1\} \cup \{b_i : 1 \leq i \leq n\}$ and $E(\Delta_n) = \{a_i a_{i+1} : 1 \leq j \leq n\} \cup \{a_i b_i : 1 \leq j \leq n\} \cup \{a_{i+1} b_i : 1 \leq i \leq n\}$ be the vertex set and the edge set of Δ_n respectively. We define a total labeling $f : V(\Delta_n) \cup E(\Delta_n) \rightarrow \{1, 2, \dots, 6n + 1\}$ as follows:

$$f(v) = \begin{cases} 3n + i & \text{if } v = a_i \text{ for } 1 \leq i \leq n + 1 \\ 5n + 2 - i & \text{if } v = b_i \text{ for } 1 \leq i \leq n \end{cases}$$

$$\begin{aligned}
 & \text{and} \\
 f(e) = & \begin{cases} i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq n \\ 3n + 1 - i & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq n \\ n + i & \text{if } e = a_{i+1} b_i \text{ for } 1 \leq i \leq n. \end{cases}
 \end{aligned}$$

To prove that f is a $C_3 - E$ -SMGL of Δ_n .

Let C_3^i for $1 \leq i \leq n$ be the subcycle of L_n with $V(C_3^i) = \{a_i : 1 \leq j \leq n\} \cup \{b_i : 1 \leq j \leq n\}$ and $E(C_3^i) = \{a_i a_{i+1} : 1 \leq j \leq n\} \cup \{a_i b_i : 1 \leq j \leq n\} \cup \{a_{i+1}, b_i : 1 \leq j \leq n\}$.

Suppose $1 \leq i \leq n$.

$$\begin{aligned}
 \text{Then } M = & \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_i) - [f(a_i a_{i+1}) + f(a_i b_i) + f(a_{i+1} b_i)] = [3n + i] + \\
 & [3n + i + 1] + [5n + 2 - i] - [i + 3n + 1 - i + n + i] = 7n + 2.
 \end{aligned}$$

The graph Δ_n is $C_3 - E$ -SMG with magic constant $7n + 2$. □

Example 2.8. The triangular snake graph Δ_6 admits C_3 -E-SMGL with magic constant $M = 44$.

Let $V(\Delta_6) = \{a_i : 1 \leq i \leq 7\} \cup \{b_i : 1 \leq i \leq 6\}$ and $E(\Delta_6) = \{a_i a_{i+1} : 1 \leq i \leq 6\} \cup \{a_i b_i : 1 \leq i \leq 6\} \cup \{a_{i+1} b_i : 1 \leq i \leq 6\}$ be the vertex set and the edge set of Δ_6 respectively. Define $f : V(\Delta_6) \cup E(\Delta_6) \rightarrow \{1, 2, \dots, 37\}$ as follows:

$$\begin{aligned}
 f(v) = & \begin{cases} 18 + i & \text{if } v = a_i \text{ for } 1 \leq i \leq 7 \\ 32 - i & \text{if } v = b_i \text{ for } 1 \leq i \leq 6 \end{cases} \\
 & \text{and} \\
 f(e) = & \begin{cases} i & \text{if } e = a_i a_{i+1} \text{ for } 1 \leq i \leq 6 \\ 19 - i & \text{if } e = a_i b_i \text{ for } 1 \leq i \leq 6 \\ 6 + i & \text{if } e = a_{i+1} b_i \text{ for } 1 \leq i \leq 6. \end{cases}
 \end{aligned}$$




Fig. 4. C_3 -E-SMGL of triangular snake Δ_6

To prove that f is a $C_3 - E$ -SMGL of Δ_6 .

Let C_3^i for $1 \leq i \leq 6$ be the subcycle of Δ_6 with $V(C_3^i) = \{a_i : 1 \leq i \leq 6\} \cup \{b_i : 1 \leq i \leq 6\}$ and $E(C_3^i) = \{a_i a_{i+1} : 1 \leq i \leq 6\} \cup \{a_i b_i : 1 \leq i \leq 6\} \cup \{a_{i+1}, b_i : 1 \leq i \leq 6\}$.

Suppose $1 \leq i \leq 6$.

$$\begin{aligned}
 \text{Then } M = & \sum_{v \in V(C_3^i)} f(v) - \sum_{e \in E(C_3^i)} f(e) = f(a_i) + f(a_{i+1}) + f(b_i) - [f(a_i a_{i+1}) + f(a_i b_i) + f(a_{i+1} b_i)] = [18 + i] + \\
 & [18 + i + 1] + [32 - i] - [19 - i + 6 + i + i] = 44.
 \end{aligned}$$

Thus the graph Δ_n is $C_3 - E$ -SMG with magic constant 44.

3 Conclusion

In this article, we have discussed C_n - E -super Magic Graceful Labeling of Some Special Graphs. Also we have given the examples related to the theorem.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Gallian JA. A Dynamic Survey of Graph Labeling. Electron. J. Combin. 2017;#DS6. 16.
- [2] Sedláček. Problem 27, in Theory of Graphs and its Applications, Proc. Symposium Smolenice. 1963;163-167. 10.
- [3] Marimuthu G, Balakrishnan M. E-super vertex magic labelings of graphs. Discrete Applied Mathematics. 2012;160(12):1766-74. 6.
- [4] Gutierrez A, Llado A. Magic coverings. Journal of combinatorial mathematics and combinatorial computing. 2005;55:43. 8.
- [5] Llado A, Moragas J. Cycle-magic graphs. Discrete Mathematics. 2007;307(23):2925-33. 14.
- [6] Rosa A. On certain valuations of the vertices of a graph. Theory of Graphs (Inter. Nat. Sym. Rome, July 1966), Gordon and Breach, N.Y. and Dunod Paris. 1967;349-355.
- [7] Golomb SW. How to number a graph, Graph Theory and Computing. Academic Press, New York. 1972;23:37. 17.
- [8] StalinKumar S, Marimuthu G. H - E -super Magic decomposition of complete bipartite graphs. Electron. Notes Discrete Math. 2015;48:297-300. 15.
- [9] Murugan S, Kumar SC. C_m - E -super magic graceful labeling of graphs. Malaya Journal of Matematik (MJM). 2019;7(4, 2019):716-9.
- [10] Andrews WS. Magic Squares and Cubes, Dover; 1960.
- [11] Ali G, Bača M, Bashir F. On super vertex-antimagic total labelings of disjoint union of paths. AKCE International Journal of Graphs and Combinatorics. 2009;6(1):11-20.
- [12] Baskar Babujee J, Vishnu Priya V. Edge bimagic labeling in graphs. Acta Ciencia Indica. 2005;XXXIM(3):741.
- [13] Bondy JA, Murty USR. Graph Theory with Applications, Elsevier, North Holland, New York; 1986.
- [14] Lin Y, Miller M. Vertex-magic total labelings of complete graphs. Bull. Inst. Combin. Appl. 2001;33:68-76.
- [15] Muthuraja NT, Selvagopal P, Jeyanthi P. Cycle-supermagic coverings and decomposition of some graphs. Amer. J. Math. Sci. Appl. 2014;2(1):83-92.
- [16] Ngurah AA, Salman AN, Susilowati L. H-supermagic labelings of graphs. Discrete Mathematics. 2010;310(8):1293-300.
- [17] Petersen J. Die Theorie der regulären Graphen. Acta Math. 1891;15:193-20.

© 2023 Sindhu and Kumar; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/98052>