



Emotional Freedom Techniques for Childhood Sexual Abuse Survivors

Maria Isabel Soriano-Lemen ^{a*} and Gina Lamzon ^a

^a *Ateneo de Davao University, Davao City, Philippines.*

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJESS/2023/v43i2936

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/98493>

Original Research Article

Received: 19/02/2023

Accepted: 21/04/2023

Published: 03/05/2023

ABSTRACT

Childhood Sexual Abuse (CSA) is a phenomenon that has affected the lives of many women and female adolescents. Eleven (11) residents of a center for abused females were included in this study. Using the case study research method of Yin, data was gathered from multiple sources. Findings were (1) emotion regulation skills were enhanced for some participants; (2) estrangement or hostility from family contributed to the prevalence of the symptoms; (3) some participants' precursor to change characteristics may have hindered therapy progress; (4) EFTs lack an emphasis on building therapeutic alliance; (5) Cognitive Behavioral Theory when used with EFTs make the sessions more helpful; (6) outside factors may have affected the worsening or reduction of symptoms; (7) there is a need to distinguish between tapping for day-to-day management of emotions and tapping on core issues; and (8) assessment, case conceptualization, and the preparation of a treatment plan play an important role, if there is to be treatment success. Recommendations include suggestions to enhance the therapeutic process, considerations that need to be made in relation to symptomatology of CSA survivors, their precursor to change characteristics, standardizing the implementation of EFTs, and integrating Cognitive Behavioral Theory with EFTs so that a more structured therapeutic approach can be made.

*Corresponding author: Email: mislemen@addu.edu.ph;

Keywords: Childhood sexual abuse; emotional freedom techniques; depression; stress; anxiety.

DEFINITIONS, ACRONYMS, ABBREVIATIONS

This part of the chapter gives definitions to the terms that have been used to ensure that unique meanings within the context of this research are defined and clarified.

Anxiety: Anxiety is described as the physical and behavioral effects of a traumatic experience.

Childhood Sexual Abuse: Childhood sexual abuse is described as sexual abuse experienced by an individual where there is no consent or where the child is unable to give consent.

Cognitive Reframing: Cognitive reframing changing the set-up statements so that cognitive shifts can occur.

Core Issues: Core issues refer to the root of an individual's problem. These are like distorted cognitions in Cognitive Behavioral Therapy.

Depression: Depression refers to symptoms of sadness, a depressed mood, lack of interest in activities that used to bring one joy, low self-worth, and the inability to engage in activities of daily living.

Emotional Helpers: Emotional helpers are qualified assistants who can implement Emotional Freedom Techniques and handle clients one-on-one or in a group setting.

Psychological Reversal: Psychological reversal in Energy Psychology underlies the failure of EFTs to work. This is considered to be an unconscious form of self-sabotage.

Reminder Statement: Reminder statement refers to shorter statements like "this remaining feeling," used while tapping on the other seven points after the Karate Chop is tapped on.

1. INTRODUCTION

The experience of Childhood Sexual Abuse (CSA) has very far-reaching effects on individuals. There are many areas in the bio psychosocial dimension that are affected because of the experience. It affects females in ways that haunt them for the rest of their lives. Not everyone gets the help that they need immediately. Some can get help years after they

experience CSA. For those who are lucky enough to get help, there are therapies available to them. However, treatment outcomes may vary because of different variables that might come into play.

It is a global concern because of how developmental trajectories may take a negative direction, causing the individual to develop psychopathological symptoms, if not psychological disorders, for as long as this experience is not processed. Around one billion children and adolescents between the ages of two to 17 have experienced physical, sexual, or emotional abuse as of 2018 [1]. These children are at higher risk of developing post-traumatic stress disorder, depression, and suicide [2]. Many countries have taken initiatives to protect victims of sexual abuse. For instance, the Center for Disease Control and Prevention in the United States have come up with strategies backed by evidence that were designed to prevent sexual violence [3]. UNICEF has been at the forefront among the organizations that are committed to ending violence against girls and women worldwide. They are doing this by encouraging policy reform in many countries all over the world [4].

In the Philippines, it has been estimated by the Philippine Commission on Women from 1999 to 2009, approximately thirteen percent (13%) of violence against women was in the form of rape. Centers have been put up by the government to help address the issues of violence against women. However, these centers are not well-funded. These centers may also need the help of licensed mental health professionals apart from government-appointed Social Workers who work with them.

In Mindanao, there are approximately four centers that cater to abused women and children [5]. However, the center where the study was conducted was not mentioned. This center houses sexually abused female adolescents and adults who are awaiting trial as well as those needing a place to live because their perpetrators are family members or prominent members in their community. This center invites Registered Psychologists to work with the residents, but they do not have the budget to pay for the services. This researcher was asked to provide therapeutic services to the residents of the facility to help equip them with coping skills

that will help them when they are eventually reintegrated into society.

Emotional Freedom Techniques have gained ground over the past decade. Many have advocated their use as a standalone therapy that is so versatile that it can resolve many, if not all, psychological symptoms. Many studies have also been conducted so that in the United States and some countries in Europe, Emotional Freedom Techniques are now considered as evidence-based therapeutic techniques to be used on PTSD symptoms of war veterans, for example. Emotional Freedom Techniques, according to research, can also be self-administered and can be added to their repertoire of coping skills that may help them in dealing with their emotional, physiological, and psychological issues. As such, this researcher wanted to explore the effectiveness of Emotional Freedom Techniques to help alleviate their symptoms of depression, anxiety, and stress. In this research it was assumed that Emotional Freedom Techniques, through the unblocking of energy points, can help improve one's psychological well-being. Through a case study research design, the researcher explored the ways through which Emotional Freedom Techniques have been effective or not effective in the relief of their symptoms.

1.1 Research Objectives

The goal of this study was to determine the effect of Emotional Freedom Techniques on psychopathological symptoms of female Childhood Sexual Abuse (CSA) survivors. Specifically, this research sought:

1. To determine the profile of the participants based on their case reports;
2. To determine the psychological and emotional state of the participants before the intervention; and
3. To determine the changes experienced by the participants after the intervention was provided to them through cross case analysis.

2. METHODOLOGY

2.1 Design

The researcher wanted to establish whether there are changes in CSA survivors when Emotional Freedom Techniques are used as a therapeutic technique. Emotional Freedom Techniques, when used as a therapeutic-technique, vary. As such, the most appropriate

way to study this is through Case Study Analysis using Case Reports of eleven (11) participants as the units of analysis. Case Study Analysis is typically used for unique programs, programs where there is difficulty in measuring program impacts, and where an understanding the program implementation in detail can help others design future intervention programs using Emotional Freedom Techniques [6].

2.2 Participants

The participants of this study were residents of the center for sexually abused females. There were eleven (11) participants who were chosen for this study. The following were the inclusion criteria that were used for the selection of the participants were (1) the participant must have been a survivor of childhood sexual abuse; (2) the participant needed to be a resident of the facility; (3) The participant lived in the facility for at least three months at the time that the research was conducted; (4) the participant was willing to participate in the study; and (5) the participant attended at least five (5) sessions of the intervention program.

2.3 Research Instruments

The following instruments were used in the study:

- The Basic Personality Inventory [7] a multidimensional psychological test to measure psychopathology and psychosocial adjustment.
- The DASS-21 [8], but the psychometrics for the tests were established by Antony, Bieling, Enns, & Swinson (1998).
- The Emotional Freedom Technique Manual [9] was used as a reference.
- The Narrative Evaluation Intervention Interview (NEII) [10] was used to qualitatively evaluate and evaluate the effectiveness of an intervention program through its process and outcome.
- The Precursors to Change matrix was also used to determine the areas in which the participants may need more help during therapy, it is used to identify which factors may be inhibiting change.

2.4 Data Procedure

The researcher used the following procedures in collecting the data for this research:

1. A letter asking for permission to conduct the research at the selected institution was

- hand-carried and given to the Center Director.
2. The participants were asked to sign the Informed Assent for if they were under 18 and the Informed Consent Form if they were over 18. The participants were assessed using the Bisaya version of the Basic Personality Inventory to establish their current psychological tendencies and pattern of answering.
 3. The EFT Intervention Program was held from February 24 to 28, 2019. The techniques used during the Interventions were the Full Basic Recipe, Sneaking Away, Tell-The-Story, the Movie Technique, and Tearless Trauma. Before each tapping round begins, the participants were asked for their Subjective Units of Distress Scale rating following the guidelines in Donn & Donn's Training Manual.
 4. An email was sent to the developer of the Narrative Evaluative Intervention Interview (NEII) to ask for permission to use the instrument as an interview guide for Key Informant Interviews to be conducted later during the conduct of the research.
 5. Prior to the commencement of the orientation, the DASS-21 was administered, scored, and interpreted. Group sessions were held in the morning and individual sessions were held in the afternoon. Two emotional helpers, who were supervised student-practitioners, were on hand to assist the researcher in the implementation of the Intervention Program. Each participant was asked what her Subjective Level of Distress score (SUDS) was at the beginning of the session. The Emotional Freedom Technique used depended on the severity of the SUDS.
 6. The researcher and the emotional helpers then took session notes that included information such as the technique used, the SUDS level prior to the tapping, and the ending SUDS level after each round of tapping.
 7. Immediately after the five-day intervention, the researcher administered the DASS-21 to track changes in their scores after the five-day intervention. The participants were then asked about their experience from the five-day intervention. Follow-ups were made 30- and 60-days after the five-day intervention using the DASS-21. Tabular information was requested from the Registered Social Worker on their sexual abuse profile.
 8. The researcher also conducted Key Informant Interviews on eleven (11)

participants using the Narrative Evaluation Intervention Interview.

9. Using the Precursor to Change matrix, the researcher determined the participants' precursor to change based on observations made by the researcher after the interviews. The researcher transcribed and translated the interviews. The researcher then prepared individual case reports of the participants. Using the individual case reports, analyses were made for each of the participants. The details of the sessions were put in a table.

2.5 Data Treatment and Analysis

The findings came from multiple sources of data. The data gathered from the participants to answer the research questions included the pretest and posttest scores, the BPI scores, data from the first big group session, session notes, and interview notes, which were integrated in the case study reports. The researcher then searched for patterns within and among the Case Reports, which were the units of analysis, by evaluating the case reports individually. Apart from the Case Reports, the Precursors to Change matrix was used to determine whether the participants were ready to change through therapy.

Additionally, Key Informant Interviews (KIIs) were conducted on the participants. The audio recordings were transcribed, translated, and analyzed for themes. Each research objective was answered and then conclusions were based on the weight of the evidence gathered. Conclusions were also made based on a cross-analysis of the case reports included in this research.

3. RESULTS

3.1 Profile of Respondents

Table 1 shows that the youngest participant was Louise, who is 13 years old while the oldest participant was Farah, who is 23 years old. The participant who has stayed at the center the longest is Mia while the participants who have stayed there for a shorter amount of time were Lorraine, Jade, and Sarah. The earliest age that a participant has been abused was Michelle at approximately two (2) years old while the oldest age that the participant has been abused were Louise and Ava at nine (9) years old. Most of the perpetrators were relatives, except for Michelle's and Mildred's perpetrators. The latter two

Table 1. Sociodemographic profile of the participant

Participant	Age	Years at Center	Age of First Abuse	Perpetrator	Case Filed	Case Resolved
Louise	13	3	9	Stepfather	Yes	Yes
Lorraine	14	0.5	8	Cousin	Yes	Yes
Jade	14	0.6	4	Uncle	Yes	Pending
Sarah	15	0.5	5	Cousin and Father	No	No
Michelle	16	2	2	Cousin, half-brother, neighbors, classmates	Yes	Pending
Sharon	16	2.3	4	Father and cousins	Yes	No data
Tabitha	18	2	6	Father	No data	No data
Mildred	19	2	7	Grandfather, brother, Unidentified Male Assailant	Yes	Brother is in prison, Grandfather is deceased
Mia	20	7	4	Brothers and Foster Father	Yes	Pending in Court of Appeals
Ava	21	5	9	Uncle	Yes	No data
Farah	23	4	4	Uncle, Cousins, and Grandfather	No data	No data

Note: Each participant was provided with a pseudonym

experienced sexual abuse by their neighbors, classmates, and an unidentified man. Among the participants, only seven (7) out of the eleven (11) have filed cases in court against their perpetrators. Only Louise and Lorraine's perpetrators are now in jail.

Table 2 shows the participants' pretest DASS-21 scores. Louise and Mia had normal pretest DASS-21 depression scores while Lorraine and Michelle had Extremely Severe scores. Also, in terms of anxiety scores, five (5) participants' scores were normal while five (5) participants' scores were in the Extremely Severe range. Finally, in terms of their Stress scores, two (2) participants were in the normal range while only one (1) participant, Michelle, was in the Extremely Severe range.

Shown in Table 3 is the elevated BPI scores of the participants. There were six (6) participants who had elevated scores in the Social Introversion dimension. There were also six (6) participants who had elevated scores in the Deviation dimension. Among the participants, only Farah had an elevated score in the Hypochondriasis dimension and only Michelle had an elevated score in the Interpersonal problem dimension. These elevated scores may also give an idea of what kind of psychological issues each participant is facing.

Based on the data gathered from the participants, the researcher filled out the Precursor to Change matrix to estimate their characteristics related to their readiness for

change. Table 4 shows the participants' Precursors Model of Change Scores.

Table 4 shows the Precursor to Change scores of the participants as well as the areas that need to be monitored for each. Michelle and Lorraine had low scores which suggest that any therapeutic changes may be limited or erratic. Further, when working with them, they need to be educated on the dimensions where their rating is low. Also, Tabitha and Mildred's scores suggested that therapeutic change is steady and noticeable. Most of the participants' scores, however, were high, suggesting that they ranged from highly motivated to inspired.

3.2 Changes Experienced by the Participants

Table 5 shows an evaluation of the movement of the DASS-21 scores of the participants. Those whose evaluations were considered effective were in normal levels by the 60-day follow-up. Those whose evaluations were considered Somewhat Effective were those participants whose scores decreased but not enough to consider the range normal. Those whose evaluations were considered Not Effective were those whose scores increased.

Themes from the Key Informant Interviews:

Table 6 shows the different themes that were extracted from the case analyses and the other data gathered by the researcher. The researcher analyzed the data and came up with themes. The main themes extracted from the data were reflected in Table 6.

Table 2. Pretest DASS-21 Scores

Participant	Age	Depression	Interpretation	Anxiety	Interpretation	Stress	Interpretation
Louise	13	8	Normal	6	Normal	18	Mild
Lorraine	14	34	Extremely Severe	22	Extremely Severe	16	Mild
Jade	14	14	Moderate	20	Extremely Severe	22	Moderate
Sarah	15	20	Moderate	16	Severe	18	Mild
Michelle	16	34	Extremely Severe	30	Extremely Severe	26	Extremely Severe
Sharon	16	18	Moderate	22	Extremely Severe	22	Moderate
Tabitha	18	12	Mild	4	Normal	6	Normal
Mildred	19	16	Moderate	6	Normal	20	Moderate
Mia	20	6	Normal	4	Normal	6	Normal
Ava	21	22	Severe	20	Extremely Severe	20	Moderate
Farah	23	23	Moderate	6	Normal	16	Mild

Table 3. Elevated BPI Scores of the Participants

Participant	Age	Elevated BPI Scores
Louise	13	None
Lorraine	14	Thinking Disorder (72)
Jade	14	Thinking Disorder (74), Social Introversion (74), Deviation (72)
Sarah	15	Self-Depreciation (87), Deviation (72)
Michelle	16	Depression (71), Interpersonal Problems (71), Alienation (79), Thinking Disorder (75), Social Introversion (81), Self-Depreciation (73), Deviation (92)
Sharon	16	Thinking Disorder (70), Social Introversion (75)
Tabitha	18	Persecutory Ideas (86), Thinking Disorder (87), Self-Depreciation (77), Deviation (92)
Mildred	18	Denial (73), Social Introversion (68)
Mia	20	Alienation (72), Persecutory Ideas (74), Thinking Disorder (83), Deviation (83)
Ava	21	Alienation (76), Persecutory Ideas (70), Thinking Disorder (92), Deviation (87)
Farah	23	Hypochondriasis (71), Persecutory Ideas (70)

Table 4. Precursors to change scores of the participants.

Participant	Age	Total Score	Scores with Lowest Ratings	Interpretation
Louise	13	23		Highly motivated to inspired. Change occurs easily. Standard approaches work well
Lorraine	14	14	Sense of Necessity (3)	Change limited or erratic: Educate client and focus on precursors with lowest ratings
Jade	14	28		Highly motivated to inspired. Change occurs easily. Standard approaches work well
Sarah	15	22		Highly motivated to inspired. Change occurs easily. Standard approaches work well
Michelle	16	10	Sense of Necessity (1), Confronting (1), Effort Toward Change (1), Hope for Change (1), Social Support (1)	Change limited or erratic: Educate client and focus on precursors with lowest ratings
Sharon	16	23		Highly motivated to inspired. Change occurs easily. Standard approaches work well
Tabitha	18	16	Sense of Necessity (2), Willingness for Anxiety (2), Awareness (2), Confronting (2), Social Support (2)	Change is steady and noticeable: Use lowest rated precursors to stay on track
Mildred	18	18	Effort Toward Change (2), Hope for Change (2), Social Support (2)	Change is steady and noticeable: Use lowest rated precursors to stay on track
Mia	20	19	Effort Toward Change (2), Hope for Change (2), Social Support (2)	Change is steady and noticeable: Use lowest rated precursors to stay on track
Ava	21	24		Highly motivated to inspired. Change occurs easily. Standard approaches work well
Farrah	23	22		Highly motivated to inspired. Change occurs easily. Standard approaches work well

Table 5. Levels of depression, anxiety, and stress of the participants

Name	Age	Depression	Anxiety	Stress
Louise	13	Effective	Somewhat Effective	Effective
Lorraine	14	Somewhat Effective	Not Effective	Not Effective
Jade	14	Effective	Somewhat Effective	Effective
Sarah	15	Not effective	Not effective	Not effective
Michelle	16	Somewhat Effective	No Change	Somewhat Effective
Sharon	16	Effective	Effective	Effective
Tabitha	18	Effective	Not Effective	Not Effective
Mildred	19	Effective	Not Effective	Effective
Mia	20	Not Effective	Not Effective	Not Effective
Ava	21	Effective	Somewhat Effective	Somewhat Effective
Farah	23	Effective	Not Effective	Effective

Table 6. Different themes extracted from the case analyses

Themes	Changes Described
Effect on emotion regulation skills of the participants	Control of emotions Release of emotions Identification of emotions Increased awareness of emotions Changing negative emotions/thoughts to positive emotions/thoughts through tapping
Improvement of cognitive functioning	Remembering tasks better Remembering lessons at school Retaining information longer
Physiological effects on participants	More relaxed after tapping Better quality of sleep because of tapping
Cathartic effect on the participants	Feeling lighter after the sessions
Reduction of Psychopathological symptoms	Reduction in temptation to self-harm No more fainting spells
Increase in Self-Awareness	Identification of weaknesses Ability to monitor thoughts to explain or rationalize good/bad behavior
Blocks to Treatment Success	Hesitance to discuss the past Inability to explore deeper into feelings to identify main issue Interpersonal problems especially with other girls in the center Trust issues prevented them from speaking up Issues with family made it difficult for some to manage their moods Wholehearted participation during the sessions Feeling uncomfortable in group sessions
Factors that contributed to treatment success	Readiness for change Willingness to explore their past Willingness to disclose personal issues Consistency in tapping
EFTs and Counseling	Counseling alone helps them come to terms with the past but tapping helps them Tapping helps them deal with their emotions There is a hesitance to speak or express oneself during counselling

4. DISCUSSION

There are findings that explain why EFTs were successful for some participants and why they were not for others. First, emotion regulation skills which may have been affected by the participants' experience of CSA were enhanced for some participants but not enough for others. Second, coming to terms with estrangement or hostility from family contributed to maintenance

of some symptoms. Third, some participants may not have been ready to change, so regardless of what therapeutic approach is used, their levels of depression, stress, and anxiety will not change. Fourth, there is a lack of emphasis for EFT practitioners to building therapeutic alliance prior to the intervention. Fifth, Cognitive Behavioral Theories, when used along with EFTs make the sessions more helpful for CSA survivors. Sixth, outside factors such as one's environment have

an effect on the maintenance or reduction of symptoms. Seventh, while tapping may work on emotions, there needs to be an emphasis that allows the client to distinguish between tapping for day-to-day management of emotions and tapping on core issues. Finally, EFTs operate in much the same way other therapies operate and can therefore be evaluated in the same way as other therapies. This makes the process of assessment and case conceptualization just as important if there is to be treatment success.

4.1 Enhancement of Emotion Regulation Skills

Emotion Regulation Skills were enhanced because of EFTs. According to Shepps, Suri, and Gross [11], the ability to regulate emotions influences the development of psychopathological symptoms. The ability to regulate emotions, according to them, involves being able to identify what they are, selecting options that help regulate them, implementing the chosen option, and monitoring the implementation overtime. In addition, Nolen-Hoeksema [12] mentioned that women tend to ruminate more than men, which partially explains the ineffectiveness of Emotional Freedom Techniques on some of the participants who have this tendency.

Childhood Sexual Abuse (CSA), according to Hebert, Langevin, and Oussaid [13], affects an individual's ability to regulate emotions. Putnam, Harris and Putnam [14] even mentioned that the experience of four or more adverse events in childhood may determine the severity of psychopathological symptoms that a CSA survivor may experience during adulthood. Some of the participants were subject to multiple abuse and neglect. Thus, it can be expected that they will experience difficulty in regulating their emotions.

In addition, it was mentioned that those who have survived early and chronic trauma, labeling and identifying emotions may pose as a problem (Briere & Scott, 2015). For example, they may not be able to distinguish anger from other emotions. In the case of Lorraine, her anger may be just homesickness or loneliness. Hence, teaching the participants the new skill of tapping to deal with intense emotions is helpful to them. This is one way they can learn to label and identify their emotions and consequently deal with them.

4.2 The Role of Estrangement and Hostility from the Family in Developing

4.2.1 Psychopathological symptoms of CSAs

Most of the participants had perpetrators who were part of the family, although not all of them are estranged from them. According to Afifi and Macmillan [15], supportive relationships and a stable environment provided by families contribute to a CSA survivor's resilience. Additionally, Domhardt, Munzer, Fegert, and Goldbeck [16] have also mentioned support from the family has an important role in ensuring that an individual is able to function normally even if they were subject to experiences of CSA. Sexual abuse that is perpetrated by family members or those who children trust, eventually results in feelings of betrayal, especially once the child begins to realize that they have been lied to [17]. It has been challenging for some participants to be so far away from their families. Lorraine worried about their parents, and this was causing her anxiety. On the other hand, Ava was very young when she moved away from her home that she is not attached to her family. In her case, there was no distress over being away from home. Research has shown that family support plays an important role in buffering the effects of Childhood Sexual Abuse. Estrangement from family thus can exacerbate effects of CSA on some of the participants. Apart from that, their perceived caregiver belief following their CSA disclosure may also play a part in the maintenance of negative emotions. Witek [18] has mentioned that perceived caregiver belief can prevent the development of psychopathological symptoms [19,20].

4.2.2 Precursors to change

One of the more important factors to ensure therapeutic success in a client's precursor to change. According to Hanna [21], looking at seven (7) variables of his Precursor to Change Model would explain why progress is made in therapy for some clients and why it would not for others. In this research, participants were rated in terms of their readiness to change based on their response to therapy and other factors. Those who scored high were able to derive the most benefits from the EFT sessions. This is a consideration that needs to be made and must be determined in the beginning of therapy. The clients' characteristics that predispose them to change strongly influence treatment progress.

4.2.3 Lack of emphasis placed on therapeutic alliance

Therapeutic alliance is composed of the bond between the client and the therapist, the agreement about the goals of therapy, and the agreement about the tasks of therapy [22]. There is hardly any literature that discusses the role of the clinician in EFT work. In the *Clinical EFT Handbook*, Frost [23] discusses how important the therapeutic or healing relationship is to ensure treatment success. However, she focuses on transference and countertransference. Donn gives an example of how to start the session. In his example, he suggested to begin the session with the client through EFTs. The session immediately begins with tapping. However, in Donn and Donn's [24] *EFT Manual*, there was a brief mention of therapeutic alliance.

The absence of a treatment plan may not allow the clinician to determine treatment length, identify the client's core issue, and work with the client's core issues. The risk here is that the client may only mention issues related to managing day-to-day emotional reactions rather than going into the details of the traumatic experience. Basically, it's like a band-aid solution to addressing what may be a bigger problem. Wampold [22] mentioned that based on his contextual model, an agreement with the client about the therapy needs to be elicited. Otherwise, the patient will not likely implement what was discussed in therapy.

4.2.4 Cognitive behavioral theories must be used along with EFTs for EFTs to work well

Although tapping the points with EFTs helped the participants become more relaxed, it was the cognitive shifts that occurred that may have had more of an effect on reducing the participants' levels of depression, anxiety, and stress. EFTs thus augment Cognitive Behavioral Therapy, and these have allowed CSA survivors to identify their core beliefs and deal with the emotions that result because of the cognitive distortions that supported their core beliefs.

A study conducted by Waite and Holder [25] mentioned that EFTs were not effective just by tapping the meridians. In fact, they were more effective in the same way traditional therapies were. Their findings suggested that EFTs were nondependent on tapping the meridian points. It worked the same way that systematic

desensitization and distraction work. In other words, a search for studies have yeti explore why EFTs work and the mechanisms through which they work because tapping alone may not guarantee therapeutic change in clients.

4.2.5 The importance of addressing core issues

While tapping has helped most of the participants in addressing their depression, anxiety, and stress symptoms, there is a need to emphasize the importance of addressing core issues. Getting to the bottom of the client's psychopathological symptoms sometimes involves learning about what is making these symptoms worse.

Assessing the client with an appropriate battery of tests that focus on the symptoms of the experience of Childhood Sexual Abuse can provide the clinician with information that is needed to address and identify the client's core issues. Apart from that, assessment can also help the clinician identify the presence of comorbid or more severe disorders, which may not be helped by Emotional Freedom Techniques. For example, three participants engaged in non-suicidal self-injury, which are indicative of a dysregulation in their stress responses, an impairment of their working models of themselves and others, and a desire to regulate emotions [26]. The experience of CSA can also manifest in different ways for different persons. However, there are commonalities among them that need to be understood and contextualized. Hence, an understanding of childhood trauma experiences and their effects on individuals is important in ensuring treatment success.

4.2.6 Specific aspects need to be explored

One of the major tasks of a therapist when using EFTs is to identify Specific Aspects of an event. These aspects for a bigger part of the picture in an emotionally charged event. According to Craig [27], specific aspects that have not been addressed can become a source of feelings that remain intense during EFT interventions. The more Specific Aspects there are within an event, the more time it will take to completely decrease emotional intensity of an event. Therapists using EFTs need to be thorough in exploring these aspects. Sometimes, during an EFT session, as one aspect is cleared, another starts to appear. This usually happens when (1) the therapist

focuses on a new detail; (2) the emotional intensity of an event using the SUDS will not decrease to zero; (3) the emotional intensity comes back; and when (4) the emotional intensity increases.

4.2.7 The need to tailor-fit EFT treatments for CSA survivors

Although there are attempts to standardize treatment for different psychological disorders such as PTSD for War Veterans, CSA survivors' needs must be considered when conceptualizing their treatment plans that are tailor-fit for them. Diagnault and Hebert [28] have studied effects on the effect of CSA on challenges faced by their participants on school and have found that students who were sexually abused experienced difficulties at school which included difficulties in their academics, behavioral difficulties such as acting out of withdrawing, presenting many psychopathological symptoms, and those who were resilient. They also suggested that there has to be differential treatment provided to these students based on their functioning.

In standardizing treatments for CSA survivors, there is also a need to recognize that they may manifest symptoms in different ways. As previously mentioned, Diagnault and Hebert identified school functioning profiles of adolescents who were sexually abused. This can be a way to assess those with CSA experiences to enhance their academic functioning. Centers that cater to CSA survivors can enhance academic functioning, behavioral problems, and interpersonal functioning by customizing treatments for this population perhaps by having these dimensions assessed and closely monitoring them.

4.2.8 Outside factors may affect effectiveness of EFT intervention

Just like with any other intervention, EFTs are effective when the client's therapeutic change and journey is supported by an environment that allows it. In the case of the participants, interpersonal relationships within the center have also acted as somewhat of a trigger to everyone living in the center. Lynskey and Fergusson [29] have found that adolescent CSA survivors are at risk for adjustment difficulties except when the nature and quality of peer and family relationships are good.

It is important to note is that cases of CSA have different symptomology and case presentations.

Everyone with an experience of sexual abuse will also have different ways of dealing with them and because of this, this will pose a challenge for clinicians who want to approach this with a standardized treatment that will apply to all cases. While treatment outcomes can be measured by success or failure in the reduction of symptoms, the approach used with everyone with childhood sexual abuse experience will likely be different for each one.

In the conduct of this study, the population was small and homogenous. However, the participants presented with diverse symptoms and CSA experiences. There is also a need to recognize that time gap between the experience of Childhood Sexual Abuse and Emotional Freedom Techniques to understand why some participants progressed faster than others.

5. CONCLUSION

In conducting this study EFTs were used techniques to help CSA survivors with their levels of depression, stress, and anxiety. There were eleven (11) participants who lived at a center for women in Mindanao, who took part in this research. The findings of this study are tentative at best. The researcher used a case study approach to answer research objectives which were to describe the participants' status prior to the intervention which were to determine the profile of the participants based on the case reports, determine the status of the participants before the intervention, and finally, to determine the changes experienced by the participants.

Numerous observations included ideas to help improve aspects of the therapeutic process and the use of EFTs during sessions. The researcher also observed that EFTs also employ Cognitive Behavioral Theories as well as principles of Energy Psychology. This researcher believes that EFTs are effective because of the use of CBT along with the tapping. Although proponents of EFT say that EFTs are now evidence-based, this researcher believes that a more structured approach in EFT sessions would enhance its effectiveness.

Results showed that that EFTs are not always effective in all cases to help reduce symptoms of depression, anxiety, and stress of CSA survivors. Many factors can affect the effectiveness of the intervention such as participant characteristics, some improvements that need to be made in

improving or enhancing the therapeutic process, and finally, the recognition that Cognitive Behavior Theory will play a major part in treatment success as well.

Taken together, these results show that EFTs can be further improved by adding more elements to the counseling process and by anchoring its practice not just on Energy Psychology but also on Cognitive Behavioral Theories. Doing so will enhance EFTs effectiveness in the longer-term.

For effective therapy for survivors of childhood sexual abuse (CSA), it is recommended that therapists consider the unique symptomatology and precursor to change characteristics of their clients. In addition, it is essential to standardize the implementation of evidence-based therapeutic approaches, such as Emotional Freedom Techniques (EFTs), to maximize therapeutic effectiveness.

To enhance the therapeutic process, therapists should also consider integrating Cognitive Behavioral Theory (CBT) with EFTs. By combining these two approaches, clients can benefit from a more structured and targeted therapeutic approach that addresses the cognitive and emotional aspects of their trauma. This integration can lead to better outcomes for clients, as they are provided with a comprehensive and effective treatment plan.

Considerations should also be made to ensure that the therapeutic approach is tailored to the unique needs of each client. This may include individualized modifications to the standard EFT protocol to accommodate the specific needs and circumstances of the client.

CONSENT

The author has requested and received approval from the respondents. Both informed consent and assent forms have been completed, signed, and documented.

ACKNOWLEDGEMENTS

The study is researcher-funded. The author acknowledges her colleagues, members of the institution, her mentor, family and friends for their support and generosity.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. World Health Organization. Guidelines for medico-legal care for victims of sexual violence; 2013 [cited Nov 29, 2018]. Available:http://www.who.int/violence_injury_prevention/resources/publications/en/guidelines_chap2.pdf.
2. Paolucci EO, Genuis ML, Violato C. A meta-analysis of published research on the effects of child sexual abuse. *J Psychol.* 2001;135(1):17-36.
3. Center for Disease Control and Prevention. Violence prevention; 2018 [cited Sep 21, 2019]. Available:<https://www.cdc.gov/violenceprevention/sexualviolence/prevention.html>
4. Ferguson H. Making home visits: creativity and the embodied practices of home visiting in social work and child protection. *Qual Soc Work.* 2018;17(1): 65-80.
5. Department of Social Welfare and Development. Centers and Institutions in the Philippines; 2019 [cited Sep 21, 2019]. Available:<https://www.dswd.gov.ph/programs/residential-and-non-residential-facilities/centers-and-institution/>
6. Balbach ED. Using case studies to do program evaluation. CA: Department of Health Services; 1999.
7. Jackson DN. The Basic Personality Inventory manual. Port Huron, MI: Research Psychologists Press; 1989.
8. Lovibond PF, Lovibond SH. The structure of negative emotional states: comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther.* 1995;33(3):335-43.
9. Donn P, Donn T. AAMET EFT Level 1 and 2 training: Emotional Freedom techniques. United Kingdom: EFT Training Centre; 2017.
10. Hasson-Ohayon I, Kravetz S, Roe D. A qualitative approach to the evaluation of interventions for persons with severe mental illness. *Psychol Serv.* 2006;3(4):262-73.
11. Sheppes G, Suri G, Gross JJ. Emotion regulation and psychopathology. *Annu Rev Clin Psychol.* 2015;11:379-405.
12. Nolen-Hoeksema S. Emotion regulation and psychopathology: The role of gender. *Annu Rev Clin Psychol.* 2012; 8:161-87.

13. Hebert M, Langevin R, Oussaid E. Cumulative childhood trauma, emotion regulation, dissociation, and behavior problems in school-aged sexual abuse victims. *J Affect Disord.* 2018;1(225): 206-312.
14. Putnam KT, Harris WW, Putnam FW. Synergistic childhood adversities and complex adult psychopathology. *J Trauma Stress.* 2013;26(4):435-42.
15. Afifi TO, MacMillan HL. Resilience following child maltreatment: a review of protective factors. *Can J Psychiatry.* 2011; 56(5):266-72.
16. Domhardt M, Munzew A, Fegert JM, Goldbeck L. Resilience in survivors of child sexual abuse: A systematic view of literature. *Trauma Violence Abuse;* 2014.
17. Finkelhor D. Early and long-term effects of child sexual abuse: An update. *Prof Psychol Res Pract.* 1990;21(5): 325-30.
18. Witek C. Resilience to posttraumatic stress symptoms in child survivors of childhood sexual abuse [cited Oct 6, 2019]. Available: <https://repository.lib.fit.edu/handle/11141/2926> [thesis/dissertation]; 2019
19. World Health Organization. *Violence Against Women;* 2017.
20. World Health Organization. *Violence against children;* June 7 2019 [cited Sep 3, 2019]. Available: <https://www.who.int/news-room/factsheets/detail/violence-against-children>
21. Hanna FJ. *Therapy with difficult clients: using the precursors model to awaken change.* Washington, DC: American Psychological Association; 2002.
22. Wampold BE. How important are the common factors in therapy? *World Psychiatry.* 2015;14(3):270-7.
23. Frost JH. Important aspects of the healing relationship in eft: bringing the unconscious into consciousness. In: Church D, Marohn S, editors. *The clinicalEFT handbook.* Fulton, CA: Energy Psychology Press. 2013; 1.
24. Donn P. Finding core issues. In: Church D, Marohn C, editors. *The clinical eft and book: A definitive resource for practitioners, scholars, clinicians, and researchers.* CA: Fulton. Energy Psychology Press. 2013;1.
25. Waite WL, Holder MD. Assessment of the emotional freedom technique: An alternative treatment for fear. *The Sci Rev of Mental Health Practice: Objective Investigations of Controversial and Unorthodox Claims in Clinical Psychology, Psychiatry and Social Work.* 2003;2(1): 20-6.
26. Fleming M, Aronson L. The relationship between non-suicidal self-injury and child maltreatment; 2019. Cornell Research Program on Self-Injury and Recovery [cited Oct 6, 2019]. Available: <http://www.selfinjury.bctr.cornell.edu/perch/resources/the-relationshipbetween-child-maltreatment-and-non-suicidal-self-injuryfinal.pdf>
27. Craig G, Craig T. Finding aspects within the EFT tapping process; 2019 [cited Oct 10, 2019]. Available: <https://www.emofree.com/efttutorial/tappingroots/aspects.html?Itemid=>
28. Diagnault IV, Hebert M. Profiles of school adaptation: Social, behavioral, and academic functioning in sexually abused girls. *Child Abuse Negl.* 2009;33(2): 102-15.
29. Lynskey MT, Fergusson DM. Factors protecting against the development of adjustment difficulties in young adults exposed to childhood sexual abuse. *Child Abuse Negl.* 1997;21(12): 1177-90.

© 2023 Soriano-Lemen and Lamzon; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/98493>