



Effect of NAA and GA₃ on Fruit Yield and Yield Attributory Traits in Phalsa (*Grewia subinaequalis* D.C.) cv. Sharbati

**Pushpendra Kumar^a, Deepak Kumar^b, Vishal Gangwar^{c*}, Om Pal^d,
Veersain^c, Avdhesh Kumar^c and A. K. Dwivedi^a**

^a Department of Fruit Science, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, Uttar Pradesh, India.

^b Department of Vegetable Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^c Department of Fruit Science, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^d Department of Horticulture, J. S. University, Shikohabad, Firozabad, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1131190

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91393>

Original Research Article

**Received 17 June 2022
Accepted 27 August 2022
Published 31 August 2022**

ABSTRACT

The experiment included four levels of NAA, 10, 20, 30, and 40 ppm, and four levels of GA₃, 10, 20, 30, and 40 ppm, administered topically and evaluated in comparison to a control (water spray). The measurements of several parameters, such as fruit set, fruit size, fruit volume, fruit weight of 100 fruits, fruit yield per plant, and fruit yield per hectare, were made. According to the experiment, GA₃ at 40 ppm enhanced fruit set, size, weight, and volume of the fruits, yield. Combining NAA+GA₃ proved effective in improving the fruit set, fruit size, yield, and quality characteristics of phalsa fruits. For higher returns in the agro-climatic conditions of Central Uttar Pradesh, phalsa growers may be advised to apply these growth regulators.

Keywords: *Phalsa; sharbati; plant growth regulators; NAA; GA₃.*

1. INTRODUCTION

Phalsa (*Grewia subinaequalis* D.C) is an important indigenous fruit by virtue of its adaptability to be grown right from kitchen garden to well organized orchard plantation and the cooling effect it is beneficial which plays considerable role in curing diseases. It belongs to the family Tiliaceae to also belongs the important fibre crop jute. It is very common in India. Phalsa is found in wild form all along the foothills of Himalayas. Commercially grown in Uttar Pradesh, West Bengal, Punjab, Madhya Pradesh and Bihar. Besides these states, it is also cultivated on a limited scale in the states of Maharashtra, Gujarat, Andhra Pradesh etc.

Phalsa berries contain 50 to 60 percent juice, 11 percent sugar, 2 to 2.5 percent acidity, 14.4 percent carbohydrate, 1.5 percent protein, 0.9 percent fat, 129 mg/100g of pulp, 89 mg phosphorus, 3.1mg iron, traces of vitamin C and 49IU vitamin [1].

“Characteristics of fruits such as length, width, pulp-stone ratio and weight of 50 fruits were increased due to foliar feeding plant growth regulators and they can also improve fruit quality, plant growth regulators improve better formation and translocation of food, hormonal signalling by gibberellins”, similar results were found by [2].

“Gibberellins decreases in acidity due to hormones application might be due to increase translocation of carbohydrates and increase metabolism due to conversion of acids to sugar. Gibberellins improved quality of fruits supported by many researches in fruit crops” such as Byas et al. [3].

“Plant growth regulators can also increase the ascorbic acid content in fruits by synthesis of catalytic activity of several enzymes and co-enzymes which are essential in ascorbic acid synthesis. So that plant growth regulators may improve physico-chemical characters like fruit length & width, weight of 50 fruits, pulp-stone ratio, ascorbic acid content, total soluble solids, sugars through better formation and translocation of carbohydrates, starch hydrolysis and early maturation phalsa fruits” by Kacha et al. [4].

“Hence plant growth regulators are very effective in improving physico-chemical attributes or quality of phalsa fruits”. Kumar [5], Garasiya et al. [6], Karole et al. [7].

“Plant growth regulators are signal molecules produced within the plant, and occur in extremely low concentrations. Hormones regulate cellular processes in targeted cells locally and, when moved to other locations, in other locations of the plant. Hormones also determine the formation of flowers, stems, leaves, the shedding of leaves, and the development and ripening of fruit. Plant hormones shape the plant, affecting seed growth, time of flowering, the sex of flowers, senescence of leaves, and fruits. Plant growth regulators improved yield of fruit plants” [2,8,9].

2. MATERIALS AND METHODS

The experiment was conducted in the Horticulture Garden of Chandra Shekhar Azad University of Agriculture and Technology, Kanpur (U.P.) during the year 2018-2019.

2.1 Experimental Design and Treatments

Sharbati Phalsa cultivar tress. That uniform, healthy and well- establishes were chosen for experiment' goal. The tress was maintained in good condition by adhering to the recommended fertilizer doses and other horticultural practices. Throughout the research, the entire orchard was kept orderly and consistently maintained. Three unique branches from each of the nine cut Ber trees were selected and utilized as a single unit (for one treatment). As a consequence, 39 units were selected on 13 Ber trees, and the experiment proceeded as planned. The experiment included thirteen treatments, each of which included foliar sprays of GA₃, NAA, control. T₁Control (00ppm water spray), T₂(10 ppm NAA), T₃(20 ppm NAA), T₄(30ppm NAA), T₅(40ppm NAA), T₆(10 ppm GA₃), T₇(20 ppm GA₃), T₈(30 ppm GA₃), T₉ (40 ppm GA₃), T₁₀(10 ppm NAA + 10 ppm GA₃), T₁₁(20 ppm NAA + 20 ppm GA₃), T₁₂(30 ppm NAA + 30 ppm GA₃), T₁₃(40 ppm NAA + 40 ppm GA₃). In the early morning with the help of knapsack sprayer with various concentration of NAA, GA₃ and their combined concentration.

2.2 Parameters of Study

2.2.1 Fruit set (%)

The number of flowers was counted on tagged shoots and percent fruit was worked out with the help of following formula:

$$\text{Fruit set (\%)} = \frac{\text{Total number of fruit set}}{\text{Total number of female flowers appeared}} \times 100$$

2.2.2 Fruit diameter (cm)

A sample of 10 berries under each treatment were randomly selected, the diameter was measured with the help of a Vernier calipers and the average diameter was expressed in cm.

2.2.3 Weight of 100 fruits (g)

Fully ripened one hundred berries from each treatment were taken by random sampling and weighed on a physical balance.

2.2.4 Volume of fruits (cm)³

Volume was determined by displacement method with the help of a measuring cylinder filled with water.

2.2.5 Yield per plant (Kg)

Fruits after each picking were weighed and total yield was calculated after adding up the value of all the harvest.

2.2.6 Yield per hectare (q/ha)

The fruits of phalsa ripe unevenly, so harvesting is done time to time during may to June. Ripe fruits are picked in every harvesting and weight recorded every time. After last picking average data was obtained as yield and average yield per hectare was determined mathematically.

3. RESULTS

The following measurements of several factors were made: Fruit set %, Fruit diameter, Fruit weight of 100 fruits, Fruit volume, Fruit yield per plant, and Fruit yield per hectare.

3.1 Fruit Set (%)

The maximum fruit set percent was observed under GA₃ @40 ppm showed 69.39% fruit set followed by GA₃ @30 ppm dose (68.38%) against the minimum of 66.55% registered under control treatment.

3.2 Fruit Diameter (cm)

The maximum fruit diameter was obtained under combined effect of NAA and GA₃ i.e. treatment T₁₃ (NAA @40 ppm + GA₃ @40 ppm) showed (0.98 cm) and the minimum fruit diameter was obtained under control treatment (T₁) i.e. 0.78 cm.

3.3 Weight of 100 Fruits (g)

The combined effect of NAA and GA₃ treatment T₁₃ (NAA @40 ppm+GA₃ @40 ppm) showed the maximum fruit weight of 100 fruits (125.62 g) followed by the treatment T₁₂ (NAA @30 ppm+GA₃ @30 ppm) i.e. 120.85cm³ and the minimum fruit weight of 100 fruits reported in control treatment (85.29 g).

3.4 Volume of Fruits (cm)³

Treatment T₉ (GA₃ @40 ppm) showed the maximum fruit volume (1.27 cm³) followed by the treatment T₁₂ (NAA @30 ppm+GA₃ @30 ppm) 1.25cm³ and the minimum fruit volume reported under control treatment (1.14cm³).

3.5 Yield per Plant (Kg)

Subjected to fruit yield per plant, the treatment T₁₃(NAA @40 ppm+GA₃ @40 ppm) showed the maximum fruit yield per plant (5.89 kg/plant) followed by the treatment T₁₂(NAA @30 ppm+GA₃ @30 ppm) i.e. 5.70 kg/plant and the minimum fruit yield per plant showed under the control treatment (3.16 kg/plant).

3.6 Yield per Hectare (q/ha)

The experimental result in terms of fruit yield per hectare showed the maximum under treatment T₁₃ (NAA @40 ppm+GA₃ @40 ppm) 78.50 kg per hectare followed by the treatment T₁₂(NAA @30 ppm+GA₃ @30 ppm) i.e. 75.98 kg/ha and the minimum fruit yield per hectare obtained under control treatment (42.10 kg/ha).

4. DISCUSSION

The blooming and fruiting of plants are greatly influenced by the quality of nourishment as well as the quantity of growth regulators present in the plant body. Both an excessive and insufficient nitrogen supply have a negative impact on blooming and fruiting; in a deficient environment, plants remain yellow with delayed fruiting and malformed fruit. Foliar sprays of NAA and GA₃ influenced the fruit set in phalsa which varied considerably. Similar results in respect of fruit set have been reported by Prasad et al. [10], Trivedi et al. [11], Balakrishnan [12] in guava, Mishra and Kriska [13] and Katiyar et al. [14] in ber and Seedkolai et al. [15] in orange. Wei Shenglin [16], however, did not find any effective result in fruit set of grapes with GA₃ treatment.

Table 1. Effect of foliar application of GA₃ and NAA on Fruit set (%), Fruit diameter(cm), Weight of 100 fruits(g), Fruit Volume (cm³), Fruit Yield per plant (kg) and Fruit Yield per hectare (q/ha) of Phalsa (*Grewia subinaequalis* D.C.) cv. Sharbati

Sr. No.	Treatments	Fruit set (%)	Fruit Diameter (cm)	Fruit Weight of 100 Fruit (g)	Fruit Volume (cm ³)	Fruit Yield per plant (kg)	Fruit Yield per hectare (q/ha)
1.	T ₁ Control (00 ppm water spray)	62.16	0.78	85.29	1.14	3.16	42.10
2.	T ₂ (10 ppm NAA)	63.40	0.84	88.63	1.19	3.45	45.93
3.	T ₃ (20 ppm NAA)	64.64	0.89	91.77	1.20	3.83	51.05
4.	T ₄ (30ppm NAA)	64.94	0.95	93.50	1.22	3.85	51.30
5.	T ₅ (40ppm NAA)	65.36	0.97	94.75	1.23	3.95	52.63
6.	T ₆ (10 ppm GA ₃)	66.55	0.86	109.61	1.18	4.25	56.64
7.	T ₇ (20 ppm GA ₃)	67.12	0.90	112.49	1.22	4.58	61.05
8.	T ₈ (30 ppm GA ₃),	68.38	0.96	116.31	1.24	4.86	64.76
9.	T ₉ (40 ppm GA ₃)	69.39	0.97	119.35	1.25	4.93	65.71
10.	T ₁₀ (10 ppm NAA + 10 ppm GA ₃)	67.64	0.88	110.12	1.20	5.45	72.61
11.	T ₁₁ (20 ppm NAA + 20 ppm GA ₃)	68.62	0.92	116.30	1.23	5.52	73.56
12.	T ₁₂ (30 ppm NAA + 30 ppm GA ₃)	69.31	0.94	120.85	1.24	5.70	75.98
13.	T ₁₃ (40 ppm NAA + 40 ppm GA ₃)	70.55	0.98	125.62	1.27	5.89	78.50
	SEm (±)	0.183	0.035	0.130	0.026	0.070	0.053
	C.D. at 5% level	0.381	0.073	0.271	0.053	0.146	0.110

GA₃ plays an important role in fruit set percent and size of fruit being responsible for hormone movement. The above functions in the present investigation under the influence of GA₃ treatment might have helped improve the fruit set percent and size of fruit of phalsa. The maximum fruit set was obtained when the plants were treated with GA₃ at 40 ppm (69.39%) concentration but it increased when NAA 40 ppm was added to it (70.55%). Similar results have been reported by Prasad et al. [17] in mango. This increase in fruit number after GA₃ application might be due to improve in plant health.

The size and weight of fruits in present trial increased by the plant growth regulators treatments. Relatively greater diameter and weight of fruits were obtained with application GA₃ at 40 ppm. The findings are in line with the reports of Rajput et al. [18] in guava, who found increase in fruit size with increase in GA₃ concentration and Biswas et al. [19] in guava observed increase in the weight of fruit with increase in GA₃ concentration. This increase in diameter and weight of fruit may be due to better supply of nutrients and photosynthates to fruits of treated plants.

Growth regulators increased fruit weight and volume in the current study, which is consistent with results published in pomegranates by Anawal et al. [20].

The application of NAA in the present investigation has given marked increase in the yield of phalsa fruits. The maximum yield per plant was recorded in GA₃ 40 ppm treated plants (4.93kg). This increase in fruit yield per plant is obviously due to the increased fruit set, greater fruit size and weight. Pandey [21] reported that improved yield of fruits the present findings are in accordance with the reports of Chandra et al. [2] in aonla and Debnath et al. [9] in phalsa. GA₃ 40 ppm treatment was found effective in increasing the yield/ha in phalsa (65.71 kg) followed by 30 ppm (69.76 kg). It was however, fortified under the combined application of NAA 40 ppm + GA₃ 40 ppm (78.50 kg). The findings agree with the reports of Pandey [21].

5. CONCLUSION

According to the results of the experiment, GA₃ at a concentration of 40 ppm improved fruit set, size, weight, and volume of fruits, and yield. Under the treatment of GA₃, a higher yield was

seen per plant and per hectare. The fruit set, fruit size, and yield attributes of phalsa fruits, however, were improved by the combined treatment of NAA+GA₃ for higher returns in Central Uttar Pradesh's agro-climatic settings, phalsa growers may be advised to use these growth regulators.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Aykroyd WR. The nutritive value of Indian foods and the planning of satisfactory diets (6thEd.). ICMR, New Delhi. 1963;10:1-126.
2. Chandra R, Manju, Rawat SS, Singh KK. (Effect of foliar application of various growth regulators on yield and quality of Aonla. cv. NA- 7. Int. J. Tropical Agr. 2015;33(3).
3. Byas PN. Studies on foliar application of micronutrients and GA₃ on yield and quality of ber(*Zizyphus mauritiana* Lamk.) fruits cv. Gola. Thesis submitted to the Master of Science in Horticulture to ND. Uni. of Agr. and Tech. Faizabad, (UP); 2014.
4. Kacha HL, Jat G, Patel SK. Performance of various plant growth regulators on yield and quality of phalsa (*Grewia asiatica* L.). Hort Flora Res. Spect. 2014;3(3):292-294.
5. Kumar S, Singh AK, Yadav AL. Effect of foliar application of GA₃, NAA, KNO₃ and Borax on fruit quality of rainy season guava cv. Lucknow-49. Plant Archives. 2010;10(1):317-319.
6. Garasiya VR, Patel NM, Bhadauria HS, Wankhede VR. Effect of plant growth regulators on quality of winterseason guava (*Psidium guajava* L.) cv. L-49 (Sardar). The Asian J. of Hort; 2013.
7. Karole B, Tiwari R. Effect of pre-harvest spray of growth regulators and urea on growth, yield and quality of ber under malwa plateau conditions, Annals of Pla. and Soil Res. 2016;18 (1):18-22.
8. Chundawat, Randhava. reported that the foliar application of GA₃, 2, 4-D and 2, 4, 5-T, increased the fruit set over control in grape fruit cv. Duncan. The best result was obtained with application 750 ppm GA₃ and 10 ppm 2, 4, 5-T; 1973.
9. Debnath A, Vanajalatha K, Momin U, Reddy M. Effect of NAA, GA₃, kinetin and ethrel on yield and quality in phalsa

- (*Grewia subinaequalis* D.C.) Asian J. Hort. 2011;6(2):474-477.
10. Prasad B, Das S, Chatterjee D, Singh UP. Effect of foliar application of urea, zinc and boron on quality of guava. Journal of Applied Biology. 2005;15(1):48-50.
 11. Trivedi N, Singh D, Bahadur V, Prasad VM, Collis JP. Effect of foliar application of zinc and boron on yield and fruit quality of (*Psidium guajava* L.). Hort Flora Research Spetrum. 2012;1(3):281-283.
 12. Balakrishnan K. Foliar spray of zinc, iron, boron and magnesium on vegetative growth, yield and quality of guava. Annals of Plant Physiology. 2000;14(2):151-153.
 13. Mishra S, Kriska B. Effect of different treatments on fruit setting in *Zizyphus jujube* Mill. Proceedings of 1st international jujube Symposium, at Agricultural University of Hebei, baoding, China. 2008: 53-54.
 14. Katiyar PN, Vikas Yadav, Singh JP. Effect of preharvest spray of NAA, GA₃ and urea of fruiting, fruit quality and yield of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. Annals of Horticulture. 2010; 3(1):92-94.
 15. Seedkolai F, Sadeghi H, Moradi H. Effects of foliar applications of nitrogen, boron and zink on auxin contents, fruit set and fruit drop in orange (*Citrus sinensis*) cv. Thompson navel. Iranian Journal of Horticultural Science. 2015;46(3): 367-378.
 16. Wei Shenglin. Effect of spraying 333 ppm and boron on the shoot growth and fruit set of Kyoho grape variety. Journal of Fruit Science. 1997;14(3):179-180.
 17. Prasad B, Ray RN, Prasad KK, Choudhary BM, Brahmachari VS. Effect of growth regulators on flowering, fruit set and fruit retention in mango. J. Res. Birsa Agriculture University. 2006;18(2): 257-260.
 18. Rajput CBS, Singh SN, Singh NPEffect of certain growth substances in guava (*Psidium guajava* L.). Haryana J. Hort. Sic. 1977;6(3-4):117-119.
 19. Biswas B, Ghosh SK, Ghosh B, Mitra SK. Effect of growth substances on fruit weight, size and quality of guava cv. L-49. Indian Agriculturist. 1988;32(4):245-248.
 20. Anawal VV, Narayanaswamy P, Ekabot SD. Effect of plant growth regulators on fruit set and yield of pomegranate cv. Bhagwa, Int. J. Scientific Res. 2015; 4(9):220-222.
 21. Pandey V. Effect of NAA, GA₃ spray on fruit retention, growth, yield and quality of ber (*Zizyphus mauritiana* Lamk.) cv. Banarasi Karaka. Orisa Journal of Horticulture. 1999;27:69-73.

© 2022 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91393>