



Effect of Nitrogen and Sulphur on Growth and Yield of Wheat (*Triticum aestivum* L.)

Koppu Ghanasai^{a*}, Rajesh Singh^{a#} and Thakur Indu^{b°}

^a Department of Agronomy, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj-211007, Uttar Pradesh, India.

^b Department of Agronomy, Naini Agricultural institute, SHUATS, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2231433

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/90935>

Original Research Article

Received 29 June 2022
Accepted 16 August 2022
Published 18 August 2022

ABSTRACT

A field experiment was conducted during *Rabi* 2022 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj (U.P). The soil of the experimental plot was sandy loam in texture, nearly neutral in soil reaction (pH 7.1), low in organic carbon (0.36 %), available N (171.48 kg/ha), available P (15.2 kg/ha) and available K (232.5 kg/ha). The experiment was laid out on Randomized Block Design with nine treatments; each replicated thrice based on one year experimentation. The treatments which are T₁: Nitrogen 80 kg/ha + Sulphur 15 kg/ha, T₂: Nitrogen 80 kg/ha + Sulphur 30 kg/ha, T₃: Nitrogen 80 kg/ha + Sulphur 45 kg/ha, T₄: Nitrogen 100 kg/ha + Sulphur 15 kg/ha, T₅: Nitrogen 100 kg/ha + Sulphur 30 kg/ha, T₆: Nitrogen 100 kg/ha + Sulphur 45 kg/ha, T₇: Nitrogen 120 kg/ha + Sulphur 15 kg/ha, T₈: Nitrogen 120 kg/ha + Sulphur 30 kg/ha, T₉: Nitrogen 120 kg/ha + Sulphur 45 kg/ha are used. The results showed that application of Nitrogen 120 kg/ha + Sulphur 45 kg/ha was recorded to significantly higher Plant height (98.13 cm), No. of Tillers/hill (10.71), Plant dry weight (17.83 g/plant), whereas significantly highest Crop growth rate (7.47 g/m²/day) was recorded with the treatment Nitrogen 80 kg/ha + Sulphur 15 kg/ha. However, significantly maximum No. of Grains/spike (47.34), Test weight (38.30 g), Grain yield (6.14 t/ha), and Straw yield (9.52 t/ha) was obtained in the treatment of Nitrogen 120 kg/ha + Sulphur 45 kg/ha as compared to other treatments.

[#] Assistant Professor;

[°] PhD scholar;

*Corresponding author: E-mail: koppughanasai@gmail.com;

Keywords: Nitrogen; sulphur; wheat; growth and yield.

1. INTRODUCTION

“Wheat (*Triticum aestivum* L.), which triggered the green revolution in the Indian subcontinent, is an important food grain providing nourishment to nearly 35 per cent of people worldwide. On a global scale, the crop is grown over an area of 211.06 million ha with a production of 566.8million tonnes. India is the second largest wheat producer in the world, next only to China and the crop has provided the fastest growth to Indian agriculture. Among cereals, wheat is next to rice in the area (24.23 million ha) and production (75.6 million tonnes)” [1]. “Wheat contributes about 60 per cent of daily protein requirement and more calories to the world diet than any other food crop. As the main staple food, wheat continues to assume greater significance in the years to come both from grain productivity and quality points of view [2,3]. Wheat is one of India’s most important cereal crops with diverse uses. Intensive cultivation has resulted in the depletion of soil nutrients to a great extent; thus, the nutrient requirement of crops has increased considerably during the last few years” [4].

“The most crucial role of N in the plant is its presence in the structure of the protein, the most important building substance from which the living material or protoplasm of every cell is made” [5,6]. In addition, “nitrogen is also found in chlorophyll, the green colouring matter of leaves. Nitrogen occupies a conspicuous place in plant metabolism. All vital processes in the plant are associated with protein, of which nitrogen is an essential constituent” [7]. “Although greater N application has produced higher yields, this is not a linear relationship (see below), and there is an optimum economic application offsetting incremental yield increase against the cost of additional N inputs, which needs to be determined for individual cultivars” (King et al., 2003). “Availability of N impacts crop development, affecting seedling establishment, tillering, canopy development as well as grain filling, all of which have the potential to affect final yield and determine the crop requirements” (King et al., 2003).

“Sulphur is another essential nutrient in all plant nutrients and a component of amino acids, which are the building block of protein. In cereal crops, sulphur ranges from 0.16-0.20%” [8,9]. “Sulphur performs many physiological functions like

synthesizing sulphur - containing amino acids, which positively improve grain quality” [10]. “Sulphur is a structural constituent of organic compounds, some uniquely synthesized by plants, providing humans and animals with essential amino acids (methionine cystine and cysteine). It is involved in chlorophyll formation and activating enzymes and is a part of vitamins biotin and thiamine (B1)” (Hegde and Sudhakara Babu, 2007). “Sulphur deficiency in crops is gradually becoming widespread due to the continuous use of sulphur-free fertilizers, high-yielding crop varieties, intensive multiple cropping systems, and higher productivity” [7].

1.1 Objectives

To evaluate on effect of nitrogen and sulphur on growth and yield of wheat and study the economics of different treatment combinations.

2. MATERIALS AND METHODS

The present examination was conducted during Rabi 2021 at Crop Research Farm, Department of Agronomy, SHUATS, Prayagraj, UP, located at 25.28°N latitude, 81.54°E longitude and 98 m altitude above the mean sea level. The experiment laid out in Randomized Block Design which consisting of nine treatments with T₁: Nitrogen 80 kg/ha + Sulphur 15 kg/ha, T₂: Nitrogen 80 kg/ha + Sulphur 30 kg/ha, T₃: Nitrogen 80 kg/ha + Sulphur 45 kg/ha, T₄: Nitrogen 100 kg/ha + Sulphur 15 kg/ha, T₅: Nitrogen 100 kg/ha + Sulphur 30 kg/ha, T₆: Nitrogen 100 kg/ha + Sulphur 45 kg/ha, T₇: Nitrogen 120 kg/ha + Sulphur 15 kg/ha, T₈: Nitrogen 120 kg/ha + Sulphur 30 kg/ha, T₉: Nitrogen 120 kg/ha + Sulphur 45 kg/ha are used. “The experimental site was uniform in topography and sandy loam in texture, nearly neutral in soil reaction (P^H 7.1), low in Organic carbon (0.38%), medium available N (225 kg ha⁻¹), higher available P (19.50 kg ha⁻¹) and medium available K (213.7 kg ha⁻¹). In the period from germination to harvest, several plant growth parameters were recorded at frequent intervals; several yield parameters were recorded after harvest. Growth parameters, plant height, and plant dry weight are recorded. The yield parameters like Seeds/spikes, Test weight, Grain yield and stover yield were recorded and statistically analyzed using analysis of variance (ANOVA) as applicable to Randomized Block Design” (Gomez K.A. and Gomez A.A. 1984).

3. RESULTS AND DISCUSSION

3.1 Growth Attributes

As can be seen in Table 1 growth parameters are summarized statistically. At 100 DAS, significantly highest plant height (98.13 cm) was recorded in the treatment with Nitrogen 120 kg/ha + Sulphur 45 kg/ha over all the other treatments. However, the treatments with application of Nitrogen 100 kg/ha + Sulphur 45 kg/ha (97.27 cm) and Nitrogen 120 kg/ha + Sulphur 30 kg/ha (97.72 cm) was statistically at par with treatment Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum plant height was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ha which is (94.33 cm). Significantly maximum number of tillers/hill (10.71) was recorded with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. However, the treatments with Nitrogen 100 kg/ha + Sulphur 45 kg/ha (10.47) and Nitrogen 120 kg/ha + Sulphur 30 kg/ha (10.56) was statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum number of tillers/hills was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ha which is (9.55). Significantly maximum plant dry weight (17.83 g/plant) was recorded with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. However, the treatments Nitrogen 100 kg/ha + Sulphur 45 kg/ha (17.22 g/plant) and Nitrogen 120 kg/ha + Sulphur 30 kg/ha (17.64 g/plant) was statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum plant dry weight was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ha (15.74). Indira Chaturvedi has applied different nitrogen levels on growth, yield and nutrient uptake of wheat (*Triticum aestivum L.*) Sharma, H. R. and Gupta, A. K. Effect of

sulphur on growth parameters and yield of some selected crops.

3.2 Yield Attributes

As can be seen in Table 2 Yield parameters are summarised statistically. At the time of harvest significantly maximum Number of Grains/spike (47.34) was recorded with Nitrogen 120 kg/ha + Sulphur 45 kg/ha over all the treatments. However, the treatments Nitrogen 100 kg/ha + Sulphur 45 kg/ha (46.61) and Nitrogen 120 kg/ha + Sulphur 30 kg/ha (47.03) was statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum number of grains/spike was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ ha which is (44.56). Significantly highest Test weight (38.30 g) was recorded with nitrogen 120 kg/ha + Sulphur 45 kg/ha over all the treatments. However, the treatments Nitrogen 100 kg/ha + Sulphur 45 kg/ha (37.67 g) and Nitrogen 120 kg/ha + Sulphur 30 kg/ha (38.00 g) which were found to be statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum test weight was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ ha (35.75). Significantly highest Grain yield (6.14 t/ha) was recorded with the treatment application of Nitrogen 120 kg/ha + Sulphur 45 kg/ha over all the treatments. However, the treatments with (5.96 t/ha) in Nitrogen 100 kg/ha + Sulphur 45 kg/ha and with (6.05 t/ha) in Nitrogen 120 kg/ha + Sulphur 30 kg/ha which were found to be statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum Grain yield was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ ha (5.04). Significantly highest Straw yield (9.52 t/ha) was recorded with the treatment

Table 1. Effect of nitrogen and sulphur on growth attributes of wheat

Treatments	Plant height (cm)	Tillers/hill	Dry weight (g/plant)
1. Nitrogen 80 kg/ha + Sulphur 15 kg/ha	94.33	9.55	15.74
2. Nitrogen 80 kg/ha + Sulphur 30 kg/ha	94.62	9.76	15.89
3. Nitrogen 80 kg/ha + Sulphur 45 kg/ha	95.85	9.97	16.64
4. Nitrogen 100 kg/ha + Sulphur 15 kg/ha	95.17	9.83	16.12
5. Nitrogen 100 kg/ha + Sulphur 30 kg/ha	96.77	10.24	16.88
6. Nitrogen 100 kg/ha + Sulphur 45 kg/ha	97.27	10.47	17.22
7. Nitrogen 120 kg/ha + Sulphur 15 kg/ha	96.29	10.10	16.37
8. Nitrogen 120 kg/ha + Sulphur 30 kg/ha	97.72	10.56	17.64
9. Nitrogen 120 kg/ha + Sulphur 45 kg/ha	98.13	10.71	17.83
F- test	S	S	S
S. Em (±)	0.31	0.11	0.20
C. D. (P = 0.05)	0.94	0.33	0.61

Table 2. Effect of nitrogen and sulphur on yield attributes and yield of wheat

Treatments	Grains/ spike	Test Weight (g)	Grain yield (t/ha)	Straw yield (t/ha)	Harvest Index (%)
1. Nitrogen 80 kg/ha + Sulphur 15 kg/ha	44.56	35.75	5.04	8.16	38.19
2. Nitrogen 80 kg/ha + Sulphur 30 kg/ha	44.87	36.05	5.17	8.33	38.30
3. Nitrogen 80 kg/ha + Sulphur 45 kg/ha	45.73	36.63	5.52	8.78	38.62
4. Nitrogen 100 kg/ha + Sulphur 15 kg/ha	45.16	36.26	5.35	8.54	38.49
5. Nitrogen 100 kg/ha + Sulphur 30 kg/ha	46.37	37.32	5.82	9.17	38.84
6. Nitrogen 100 kg/ha + Sulphur 45 kg/ha	46.61	37.67	5.96	9.23	39.24
7. Nitrogen 120 kg/ha + Sulphur 15 kg/ha	45.96	37.05	5.66	8.89	38.87
8. Nitrogen 120 kg/ha + Sulphur 30 kg/ha	47.03	38.00	6.05	9.34	39.30
9. Nitrogen 120 kg/ha + Sulphur 45 kg/ha	47.34	38.30	6.14	9.52	39.21
F test	S	S	S	S	NS
S. Em (\pm)	0.25	0.26	0.07	0.10	0.29
CD (P = 0.05)	0.75	0.76	0.21	0.31	-

application of Nitrogen 120 kg/ha + Sulphur 45 kg/ha over all the treatments. However, the treatments with (9.23 t/ha) in Nitrogen 100 kg/ha + Sulphur 45 kg/ha and with (9.34 t/ha) in Nitrogen 120 kg/ha + Sulphur 30 kg/ha which were found to be statistically at par with Nitrogen 120 kg/ha + Sulphur 45 kg/ha. The minimum straw yield was recorded in the treatment combination of Nitrogen 80 kg/ha + Sulphur 15 kg/ha (8.16). There was no significant difference between the treatments. However, the highest harvest index (39.30%) was recorded with the treatment application of Nitrogen 120 kg/ha + Sulphur 45 kg/ha. In contrast, the lowest harvest index (38.19%) was recorded with nitrogen 80 kg/ha + Sulphur 15 kg/ha. Test weight, a significant yield determining component, is a genetic character and is least influenced by the environment [11-13]. This might be due to the higher conservation of light energy into chemical energy and its subsequent translocation from source to sink. Similar findings were observed by Saha et al. [14].

4. CONCLUSION

It is concluded that the application of Nitrogen 120 kg/ha + Sulphur 45 kg/ha recorded a significantly higher Grain yield (6.14 t/ha) compared to other treatments since the findings are based on the research done in one season.

ACKNOWLEDGEMENT

I thank my advisor Dr. Rajesh Singh and all the faculty members of Department of the Agronomy, Naini Agricultural Institute, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj -211007, Uttar Pradesh. For

providing us with essential facilities to undertake the studies.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Jagshoran, Sharma RK, Tripathi SC. New varieties and production. The Hindu, Survey of Indian Agriculture. 2004;33-35.
- Degra ML, Pareek BL, Shivran RK. Effect of sulphur and integrated weed management on productivity and quality of Indian mustard (*Brassica juncea*) and succeeding fodder pearl millet. Research on Crops. 2008;9(3):573-577.
- Indira Chaturvedi. Effects of different nitrogen levels on growth, yield and nutrient uptake of wheat (*Triticum aestivum* L.). International Journal of agricultural Sciences. 2006;2(2):372-374.
- Pandey GP, Khandkar UR, Tiwari SC, aNarendra Kumawat. Productivity, profitability of wheat and soil fertility as influenced by different levels of nitrogen under Sodic vertisols. International Journal of Chemical Studies. 2018;6(4):3292-3295.
- Bhatta RD, Amgain LP, Subedi R, Kandel BP. Assessment of productivity and profitability of wheat using Nutrient Expert. Heliyon. 2020;6:04144.
- Dadhich LK, Gupta AK. Plant morphology, yield, nutrient content and uptake of fodder pearl millet as influenced by S, Zn supply and its intercropping with cowpea. Annals

- of Agriculture Bio Research. 2004;9(2): 183-188.
7. Singh G, Santosh Kumar, Gurmeet Singh, Navdeep Singh. Effect of integrated nutrient management on nutrient uptake and grain yield of wheat (*Triticum aestivum* L.) under irrigated conditions. Journal of Pharmacognosy and Phytochemistry. 2019;8(1):1077-1080.
 8. Kaur R, Kumar S, Kaur R, Kaur J. Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.) under irrigated conditions. International Journal of Chemical Studies. 2018;6(4): 1800-1803.
 9. Maurya RN, Singh UP, Sunil Kumar, Yadav AC, Yadav RA. Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.). International Journal of Chemical Studies. 2019;7(1):770-773.
 10. Chaudhary PD, Jat RS, Sharma HS. Interaction effect of phosphorus, sulphur and PSB inoculation on growth, yield and nutrient uptake of wheat. Annals of Agricultural Research. 2003;24(1):12-16.
 11. Sharma HR, Gupta AK. Effect of sulphur on growth parameters and yield of some selected crops. Annals of Agricultural Research. 2003;24(1):136-138.
 12. Singh B, Kamlesh Kumar, Kaur M. Effect of nitrogen and zinc on yield and economics of winter maize (*Zea mays* L.) Under irrigated condition of Punjab. Journal of Pharmacognosy and Phytochemistry. 2019;SP1:574-576.
 13. Yadav KS, Nand H. Site-specific nutrient management for optimal food grain production in Haryana. Better Crops with Plant Food. 2004;88(2):21-23.
 14. Saha B, Parimal Panda, Patra PS, Ranajit Panda, Kundu A, Singha Roy AK, Nabakishor Mahato. Effect of Different levels of Nitrogen on Growth and Yield of Rice (*Oryza sativa* L.) Cultivars under Terai-agro Climatic Situation. International Journal of Current Microbiology and Applied Sciences. 2017;6(7):2408-2418.

© 2022 Ghanasai et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/90935>