



Quantum Probabilities and the Born Rule in the Intuitionistic Interpretation of Quantum Mechanics

Arkady Bolotin^{1*}

¹ Ben-Gurion University of the Negev, Beersheba, Israel.

Author's contribution

The sole author designed, analyzed and interpreted and prepared the manuscript

Article Information

DOI: 10.9734/PSIJ/2017/31668

Editor(s):

- (1) Daniel Beysens, OPUR International Organization for Dew Utilization, France.
(2) Stefano Moretti, School of Physics Astronomy, University of Southampton, UK.

Reviewers:

- (1) A. Ayeshamariam, Khadir Mohideen College, India.
(2) Manuel Malaver de la Fuente, Maritime University of the Caribbean, Venezuela.
(3) K. Muralidhar, National Defence Academy, Pune, India.
(4) Balwant Singh Rajput, Kumaun University, India.
Complete Peer review History: <http://www.sciencedomain.org/review-history/18303>

Received: 19th January 2017

Accepted: 9th March 2017

Published: 23rd March 2017

Original Research Article

ABSTRACT

This paper presents a novel explanation of the cause of quantum probabilities and the Born rule based on the intuitionistic interpretation of quantum mechanics where propositions obey constructive (intuitionistic) logic. The use of constructive logic makes it possible (through a replacement of the concept of truth with the concept of constructive probability) to abandon the law of excluded middle in the intuitionistic interpretation.

Keywords: Born rule; quantum probabilities; intuitionistic interpretation; quantum mechanics.

*Corresponding author: E-mail: arkadyv@bgu.ac.il;

1 INTRODUCTION

Whence comes indeterminism in physics? Really, assuming that the Schrödinger dynamics is universally valid, i.e., that each process in physics is governed by the (linear differential) Schrödinger equation and so is continuous, causal, and reversible, how is that there are processes involved in measurement that are indeterministic, i.e., whose outcomes are probabilities? It is no exaggeration to say that the problem this question relates to is the most difficult in the foundation of quantum mechanics.

Under the Copenhagen interpretation, quantum probabilities enter physics through the Born rule, which cannot be investigated as it is a postulate of the theory (the same one as the Schrödinger equation is). However, the problem with such an *ab initio* postulation is not that it does not offer any deeper understanding of how probabilities come into existence. The real problem here is that there is nothing inherent in the Copenhagen interpretation that can either reject the mutual application of the Schrödinger equation and the Born rule to a system of interest or select one postulate over another [1].

Hence, attempts to understand the cause of indeterminism in quantum physics are being made beyond the Copenhagen interpretation.

For example, in the relative state (also referred to as many-worlds) interpretation (henceforward RS/MWI for short), the behavior of a rational agent in quantum-mechanical situations is thought to provide a natural account of the probability concept [2, 3, 4, 5, 6]. As it is argued in the quantum-mechanical version of decision theory, being strongly constrained in their behavior the rational agents would quantify their subjective uncertainty in the face of the wave function “splitting” (or “branching”) by the use of probability given by the Born rule.

But then the fact that the RS/MWI contains neither element nor quality nor attribute related to a notion of *belief* makes a doubt if there be

a way to derive probabilities i.e., *degrees of beliefs* from such an interpretation.¹ What is more, even if this notion had been added to the RS/MWI, it still would be flatly inconsistent with the universally applicable (deterministic) Schrödinger dynamics and the splitting picture [9]. Such an argumentation may explain why no decisive conclusion on the possibility of emergent probabilities and randomness in the RS/MWI has been reached yet.

Meanwhile, the decoherence program is seeking the solution to the problem of the origin of quantum probability by including the environment but without relying on the key elements of decoherence that presume the Born rule and would thus render the contention circular [10, 11]. For example, as it has been proposed in the decoherence program, the Born rule is originated from the environment-assisted invariance (in other words, from a symmetry of composite quantum states) [12, 13, 14].

But then again, it is hard to see how probabilities can emerge within the framework, in which only unitary evolutions (without collapse mechanism) are allowed and thus there do not exist measurement outcomes, that is, *the bearers of probabilities*.²

So, the intention of this paper is to give explanation of the cause of quantum probabilities and the Born rule using a completely different approach. The key idea is to employ the intuitionistic interpretation of quantum mechanics [17] in which propositions concerning quantum-mechanical situations obey constructive (intuitionistic) logic. As the objects of belief that bear logical values (‘true’ \top and ‘false’ \perp) [18], propositions let one pave the way for the entrance of a probabilistic concept into quantum theory. Also, the use of intuitionistic logic makes it possible (through a replacement of the concept of truth with the concept of constructive provability) for the assumption of the universal validity of the Schrödinger dynamics to *become weaker*. Such a deregulation allows one to drop *the law of excluded middle* in the intuitionistic interpretation

¹The concept of an agent’s degree of confidence, a *graded belief*, is one of the main concepts of probability. See papers [7, 8] that analyze in depth this and other concepts of probability.

²According to the propensity, subjective, and logical interpretations of probability, measurement outcomes (or events) are the bearers of probabilities; see, for example [15, 16].

so that the last-named does not “fall victim to Schrödinger’s cat and the like” [19].

The rest of the paper is structured as follows: The essence of the points maintained in the paper is put in the next Section 2, while the Section 3 concludes the arguments of this paper.

2 HOW IRREDUCIBLE PROBABILITIES APPEAR IN THE INTUITIONISTIC INTERPRETATION

Assume that a (typical microscopic) system S , whose pure quantum states are determined by

the eigenstates $|a_1\rangle$ and $|a_2\rangle$ of the observable A (that takes on only two possible values a_1 and a_2), interacts for the duration of the measurement t with a (typical macroscopic) apparatus M designed to measure (observe) A .

Suppose that the state of the system S is a superposition $|\psi\rangle = c_1|a_1\rangle + c_2|a_2\rangle$ where coefficients c_1 and c_2 meet with the normalization condition $c_1^2 + c_2^2 = 1$ (whose reason will be explained later). In this case, the sample space of the measurement – i.e., the set of all possible outcomes of the measurement of the observable A – can be defined as $\Omega \equiv \{a_1, a_2\}$. Let us evaluate the following proposition:³

$$(\{a_1, a_2\}) \equiv ((\{a_1\}) \oplus (\{a_2\})) \equiv (((\{a_1\}) \vee (\{a_2\})) \wedge (\neg(\{a_1\}) \vee \neg(\{a_2\}))) \quad , \quad (2.1)$$

where \oplus stands for the logical operation of exclusive disjunction (corresponding to the construct “either ... or”) that outputs ‘true’ only when its inputs – i.e., the propositions of events $\{a_1\}$ and $\{a_2\}$ (subsets of $\{a_1, a_2\}$) – differ, namely, $(\{a_1\}) \neq (\{a_2\})$. The proposition $(\{a_1, a_2\})$ is logically equivalent to the assertion that there is a final (i.e., at the moment t in the last part of the measurement) state $|\Psi_t\rangle$ of the composite system $S + M$ in which the observable A has a definite value, in other words, there is an event which contains only one true outcome – *either* $\{a_1\}$ *or* $\{a_2\}$ in the sample space $\{a_1, a_2\}$ of the measurement.

Consider the first part of the compound expression (2.1), that is, the proposition $((\{a_1\}) \vee (\{a_2\}))$. In view of the fact that this proposition contains the logical constant \vee , constructive logic [20] (adopted in the intuitionistic interpretation of quantum mechanics presented here) requires that if this proposition is asserted to be true, then *witness* must be given, which chooses $(\{a_1\})$ or $(\{a_2\})$ and provides (an explicit, constructive) proof for it.⁴ What counts as “witness” is open to interpretation though. In the said case, witness can be understood as an outcome of either *an actual measurement of A* or a decision problem that determines the logical value of the proposition $((\{a_1\}) \vee (\{a_2\}))$ *before the actual measurement of A*.

This decision problem can be defined as the following propositional formula:

$$((\{a_1\}) \vee (\{a_2\})) \equiv (|\Psi_t\rangle = |a_1\rangle|M_1 \vee |\Psi_t\rangle = |a_2\rangle|M_2) \Big| P(|\Psi_t\rangle) \quad , \quad (2.2)$$

³In this paper, propositions and statements are used interchangeably and denoted (if not stated otherwise) as enclosed in parentheses expressions (\cdot) which are capable of being true or false.

⁴Another logical constant whose interpretation requires constructive proof is \exists .

where $|M_1\rangle$ and $|M_2\rangle$ represent mutually orthogonal quantum states of M , both orthogonal to $|M_0\rangle$ (which is the initial – i.e., at the moment $t = 0$ preceding the measurement – quantum state of M , so-called *ready state*), corresponding to different macroscopic configurations of M similar to different positions of a pointer along a scale, while $P(|\Psi_t\rangle)$ represents a statement – i.e., a predicate – that may be true or false depending on the particular final state $|\Psi_t\rangle$ of the composite system $S + M$.

By contrast, when the state of the system S is just one of the eigenstates $|a_n\rangle$ of the observable A , that is, $|\psi\rangle = |a_n\rangle$ (where $n \in \{1, 2\}$), the sample space of the measurement correspondingly consists of a single element, i.e., $\Omega \equiv \{a_n\}$. In the given case, a witness for the proposition of the elementary event ($\{a_n\}$) (asserting that in the final state $|\Psi_t\rangle = |a_n\rangle|M_n\rangle$ the observable A has the definite value a_n) is

required to do nothing but only exist if ($\{a_n\}$) is true. Therefore, the proposition ($\{a_n\}$) can be treated in accordance with the laws of classical logic [21]. In particular, the following propositional expression

$$(\{a_n\}) \equiv (|a_n\rangle|M_n\rangle) = \top \quad (2.3)$$

(where symbol \equiv positioned after the parentheses defines the notion of evaluation) can be taken as a postulate assuming *the eigenvector-eigenvalue link* the perfect correlation between the initial state $|a_n\rangle$ of the system S and the final state $|M_n\rangle$ of the apparatus M .⁵

Let us return to the propositional formula (2.2). Since the quantum evolution of the system $S + M$ is given as $|\Psi_t\rangle = \hat{U}(t)|\psi\rangle|M_0\rangle$, where $\hat{U}(t)$ stands for the time evolution operator, namely, $\hat{U}(t) = \exp(-itH_{S+M}/\hbar)$, the predicate $P(|\Psi_t\rangle)$ is defined such

$$P(|\Psi_t\rangle) \equiv \left((|\Psi_t\rangle = \hat{U}(t)|\Psi_0\rangle) \wedge (|\psi\rangle = c_1|a_1\rangle + c_2|a_2\rangle) \wedge (\exists|M_0\rangle \in \mathcal{H} P(|M_0\rangle)) \right) \quad (2.4)$$

that it includes the statement ($|\Psi_t\rangle = \dots$) asserting the validity of the Schrödinger dynamics for the measurement process, the statement ($|\psi\rangle = \dots$) declaring the validity of the quantum superposition principle for the system S , and the statement ($\exists|M_0\rangle \dots$) affirming the existence of the ready state of the apparatus M as a quantum state vector $|M_0\rangle$, in which \mathcal{H} denotes an abstract (separable, infinite-dimensional) Hilbert space and $P(|M_0\rangle)$ stands for the propositional function of the vector $|M_0\rangle$.

As it follows, to assign a logical value to the proposition ($(\{a_1\}) \vee (\{a_2\})$) in advance of the measurement, the predicate $P(|\Psi_t\rangle)$ must be calculated. But to do so, a witness is required that can provide computational evidence supporting the statement ($\exists|M_0\rangle \dots$) since this is the one in the formula (2.4) that involves the existential quantification \exists .

Were the state vector $|M_0\rangle$ to exist, it would be a vector of the Hilbert space $\mathcal{H}_M \subseteq \mathcal{H}$ of the apparatus M (as much as the state vectors $|M_1\rangle$ and $|M_2\rangle$ would be). From another side, the eigenvectors $|M_n\rangle$ of the Hamiltonian operator H_M corresponding to the total energy of the apparatus M must provide an orthonormal basis $\{|M_n\rangle\}$ for the Hilbert space \mathcal{H}_M they span. For this reason, let us consider *the decision problem of the Schrödinger equation* $\Pi(H_M)$ for the Hamiltonian operator H_M

$$\Pi(H_M) \equiv (\{|M_n\rangle\} \neq \emptyset) \quad (2.5)$$

whose output is taken to be true if the solution set $\{|M_n\rangle\}$

⁵Here it is also assumed the absence of degeneracy meaning there is only one eigenstate for each eigenvalue.

$$\{|M_n\rangle\} \equiv \left\{ |M_n\rangle \in \mathcal{H} \mid i\hbar \frac{\partial}{\partial t} |M_n\rangle = H_M |M_n\rangle \right\} \quad (2.6)$$

(i.e., the set of all vectors $|M_n\rangle$ for which the Schrödinger equation with H_M holds) is not empty and ‘false’ otherwise.

Let us consider the following claim

$$\Pi(H_C) = \top \quad (2.10)$$

Most generally, the decision problem of the Schrödinger equation can be presented in the form

$$\Pi(H_C) \equiv (\exists |u\rangle \in \mathcal{H} P(|u\rangle, H_C)) \quad (2.7)$$

It is clear that this claim (logically equivalent to the assertion of the universal validity of the Schrödinger dynamics) could be proved if the Schrödinger equation was capable of being exactly solved by a generic algorithm, i.e., for the Hamiltonian operator H_C .

where the predicate $P(|u\rangle, H_C)$ encloses the general time-dependent Schrödinger equation

$$P(|u\rangle, H_C) \equiv \left(i\hbar \frac{\partial}{\partial t} |u\rangle = H_C |u\rangle \right) \quad (2.8)$$

However, such an algorithm does not exist (at least as it is known in the present state of our knowledge [22]). To be exact, the decision problem $\Pi(H_C)$ is known to be undecidable, i.e., there does not exist a single (generic) method that can in a finite number of steps correctly solve this problem for any allowable Hamiltonian operator H (even though methods of finding the solution set $\{|u\rangle\}$ of the Schrödinger equation with some particular H are known).

in which H_C symbolizes a completely arbitrary Hamiltonian operator (which means that the term H_C is free for substitution for any element in the set of all allowable Hamiltonian operators H) and the vector $|u\rangle$ represents the exact solution to this equation.

Indeed, as it is argued in [17], the decision problem of the Schrödinger equation is parallel to the general spectral gap problem that asks whether a given Hamiltonian operator H has a spectral gap (i.e. the energy difference between the ground state and the first excited state of the system in the thermodynamic limit). To be sure, if there were a generic algorithm capable of obtaining the solution set $\{|u\rangle\}$ of the Schrödinger equation for all allowable Hamiltonian operators H , then such an algorithm would be able to answer not only the question whether this solution set $\{|u\rangle\}$ is empty or not but also the question whether the spectrum of the eigenvalues corresponding to $\{|u\rangle\}$ is discrete and gapped or continuous and gapless. Yet, according to the result of the paper [23], the general spectral gap problem is undecidable. One can infer from this conclusion that the generic algorithm for solving the Schrödinger equation with an arbitrary H_C does not exist and consequently the decision problem $\Pi(H_C)$ is undecidable.

Contrary to microsystems, a typical macroscopic object with its uncontrolled and unlimited degrees of freedom cannot be assigned a specific (i.e., explicit, detailed and unambiguous) Hamiltonian operator H_M . This means that the truthfulness of the statement $(\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle))$ would be secured if and only if the solution set $\{|u\rangle\}$ of the Schrödinger equation was *in no case* empty, that is, the decision problem of this equation had to be *universally true* (i.e., had the truth output for all physical systems), namely,

$$\Pi(H_C) \leftrightarrow (\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle)) \quad (2.9)$$

(where – in line with the notion of constructiveness – the symbol \leftrightarrow may be viewed as an abbreviation of “can be replaced in a proof with”).

In order to ensure the general undecidability of this problem, in the intuitionistic interpretation of quantum mechanics the declaration of the universal validity of the Schrödinger dynamics is *weakened* (in comparison with other interpretations that are based on classical or quantum logic). Specifically, it is asserted that unless one has a proof that for the particular system $\Pi(H) = \top$, the Schrödinger dynamics can be considered only being *possibly valid* for this system. Symbolically, this can be written down as the following entailment

$$\left(\exists H \Pi(H) = \top \right) \vdash \left(\diamond \Pi(H_C) = \top \right) \quad , \quad (2.11)$$

where \diamond stands for the modal operator of possibility [24]. This entailment assigns the truth value \top to the proposition “it is possible that $\Pi(H_C)$ ” as a logical consequence of the fact that for some Hamiltonian operators H the solution set of the Schrödinger equation can be explicitly demonstrated⁶.

By the axioms of seriality and reflexivity of modal logic, i.e., $\square \Pi(H_C) \rightarrow \diamond \Pi(H_C)$ and $\square \Pi(H_C) \rightarrow \Pi(H_C)$ respectively (where \square is the operator of necessity), the equality $\diamond \Pi(H_C) = \top$ suggests $\Pi(H_C) = \top$ but it equally may be $\Pi(H_C) = \perp$. This means that the entailment (11) declares *contingency* of the decision problem $\Pi(H_C)$ (in other words, the outcome of $\Pi(H_C)$ is asserted to be uncertain). As a result, from the formula (2.9) one infers the following evaluation

$$\left(\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle) \right) = \{ \} \quad , \quad (2.12)$$

where symbol $\{ \}$ represents the lack of the logical values. The meaning of the evaluation (2.12) is that the statement asserting the existence of the state vector $|M_0\rangle$ cannot be decided computationally, i.e., by means of the Schrödinger equation.

This implies that the predicate $P(|\Psi_t\rangle)$ presented in the formula (2.4) simply *cannot be known*: Its truth or falsehood cannot be analyzed (computed). The unknowingness of the predicate $P(|\Psi_t\rangle)$ in turn implies that it is impossible to assign a definite logical value to $(\{a_1\} \vee \{a_2\})$ – and so to the proposition $\{a_1, a_2\}$ – in advance of the actual measurement of the observable A . In other words, because of the undecidability of the decision problem $\Pi(H_C)$, the logical values of the propositions of elementary events $\{a_1\}$ and $\{a_2\}$ cannot be decided (and thus exist) ahead of the measurement.

In passing, let us note that *postulating the universal validity* of the Schrödinger dynamics and in this way the existence of the ready state of the apparatus M as a state vector $|M_0\rangle$, one gets a contradiction known as *the quantum measurement problem* (or, to be more exact, the problem of definite outcomes). Truly, when the both statements $(\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle))$ and $(\psi = c_1|a_1\rangle + c_2|a_2\rangle)$ are presumed to be true, the statement asserting the validity of the Schrödinger dynamics for the measurement process, $(|\Psi_t\rangle = \hat{U}(t)|\Psi_0\rangle)$, cannot be true: The linearity of the Schrödinger equation entails that in this case the initial state of the composite system $S + M$ evolves into the following final state

$$(c_1|a_1\rangle + c_2|a_2\rangle)|M_0\rangle \xrightarrow{t} c_1|a_1\rangle|M_1\rangle + c_2|a_2\rangle|M_2\rangle \quad , \quad (2.13)$$

⁶In contrast to the intuitionistic interpretation, the RS/MWI and the decoherence program infer the universal validity of the Schrödinger dynamics from the same fact, i.e., $(\exists H \Pi(H) = \top) \vdash (\Pi(H_C) = \top)$. Indeed, as it is stated for example in [25], since “there is satisfactory and often excellent evidence” that the quantum mechanical (QM) framework “is quantitatively valid” and at the same time “there is, at least at present, no positive experimental evidence that it is not valid in other regions where it has not been directly tested”, then “the principle of Occam’s razor would certainly suggest that the intellectually economical attitude is to assume that the general conceptual scheme embodied in QM is in fact valid for the whole of the physical universe without restriction”.

in which the symbol + representing the linear superposition cannot be replaced by the coordinating conjunction “or” such that $|\Psi_t\rangle = |a_1\rangle|M_1\rangle$ or $|\Psi_t\rangle = |a_2\rangle|M_2\rangle$ (meaning the inclusive or exclusive disjunction). So, without supplying an additional postulate (for example, the wave-packet reduction postulate) or giving a suitable interpretation of the superposition $c_1|a_1\rangle|M_1\rangle + c_2|a_2\rangle|M_2\rangle$, it is impossible to explain the definite pointer position $|M_n\rangle$ corresponding to the value a_n , that is, the logical value ‘true’ of the proposition $(\{a_1, a_2\})$, which is always perceived as the outcome of the actual measurement of the observable A .

Let us now evaluate the intuitionistic (Heyting) negation of the predicate $P(|\Psi_t\rangle)$:

$$\neg P(|\Psi_t\rangle) \equiv \neg \left((|\Psi_t\rangle \neq \hat{U}(t)|\Psi_0\rangle) \vee (\psi \neq c_1|a_1\rangle + c_2|a_2\rangle) \vee \neg (\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle)) \right) \quad (2.14)$$

From the definition of the decision problem $\Pi(H_C)$ it follows that its negation, namely, $\neg\Pi(H_C) \equiv (\forall |u\rangle \in \mathcal{H} \neg P(|u\rangle, H_C))$, is not constructively provable: In order to produce direct evidence of the truthfulness of this negation, one would have to demonstrate falsity of the Schrödinger equation, that is, $\neg P(|u\rangle, H_C) \equiv (i\hbar \frac{\partial}{\partial t} |u\rangle \neq H_C |u\rangle) = \top$, for all vectors $|u\rangle$ in the abstract infinite dimensional Hilbert space \mathcal{H} .

Instead, let us consider the double negation introduction

$$\neg\Pi(H_C) \leftrightarrow \neg(\neg\neg\Pi(H_C)) \quad (2.15)$$

which is a theorem in constructive logic. Since to prove $\neg\neg\Pi(H_C)$ intends to show that the solvability of the general Schrödinger equation would not be contradictory, the double negation $\neg\neg\Pi(H_C)$ has ‘true’ output when there is no evidence against universal validity of Schrödinger dynamics. Otherwise stated, $\neg\neg\Pi(H_C) = \top$ holds if $\Pi(H_C)$ cannot be falsified or it is possibly true.

As it has been already noticed, the truthfulness of $\neg\neg\Pi(H_C)$ (i.e., $\diamond\Pi(H_C) = \top$) can be inferred from the existence of any known exactly solvable quantum model. Thus, from (2.15) it follows that

$$\neg\Pi(H_C) = \perp \quad (2.16)$$

The last inference means that in the intuitionistic interpretation *not-Schrödinger dynamics* is considered *impossible* (i.e., false and necessarily false, that is, $\Box\neg\Pi(H_C) = \perp$).⁷

Taking into account the negation of the equivalence (2.9), this impossibility yields

$$(\neg\Pi(H_C) = \perp) \vdash (\neg(\exists |M_0\rangle \in \mathcal{H} P(|M_0\rangle)) = \perp) \quad (2.17)$$

At the same time, in view of the deduction $(\neg\Pi(H_C) = \perp) \vdash (\neg P(|u_C\rangle, H_C) = \perp)$, where $|u_C\rangle$ denotes a completely arbitrary vector in \mathcal{H} , and the equivalence (holding true for small t or if the Hamiltonian operator H_{S+M} does not depend on t)

$$(|\Psi_t\rangle = \hat{U}(t)|\Psi_0\rangle) \leftrightarrow P(|\Psi\rangle, H_{S+M}) \quad (2.18)$$

where the propositional function $P(|\Psi\rangle, H_{S+M})$ encloses the Schrödinger equation for $|\Psi\rangle$ and H_{S+M} , one infers that

$$(\neg P(|u_C\rangle, H_C) = \perp) \vdash ((|\Psi_t\rangle \neq \hat{U}(t)|\Psi_0\rangle) = \perp) \quad (2.19)$$

The expressions (2.17) and (2.19) together with the evaluation $(\psi \neq c_1|a_1\rangle + c_2|a_2\rangle) = \perp$ (which

⁷It also generates $\Pi(H_C) \vee \neg\Pi(H_C) \neq \top$, which means that in the presented intuitionistic interpretation of quantum mechanics, the law of excluded middle is not admitted as an axiom.

is trivial if the system S is microscopic⁸), result in that the negation $\neg P(|\Psi_t\rangle)$ must be false for any predicate variable $|\Psi_t\rangle$.

As it follows, the negation of the formula (2.2) must be 'false': $(\neg(\{a_1\}) \wedge \neg(\{a_2\})) = \perp$.

Furthermore, since the predicate $P(|\Psi_t\rangle)$ of the propositional formula for the second part of the compound expression (2.1), namely, $(\neg(\{a_1\}) \vee \neg(\{a_2\}))$, is the same as one presented in the formula (2.4), that is,

$$\begin{aligned} & (\neg(\{a_1\}) \vee \neg(\{a_2\})) \\ \equiv & (|\Psi_t\rangle \neq |a_1\rangle|M_1) \vee (|\Psi_t\rangle \neq |a_2\rangle|M_2) \Big| P(|\Psi_t\rangle) \end{aligned} \quad (2.20)$$

the negation of this part must be 'false' too: $(\neg\neg(\{a_1\}) \wedge \neg\neg(\{a_2\})) = \perp$.

So, if the propositions $(\{a_n\})$ can be assigned logical values and *for that reason* double negation elimination $\neg\neg(\{a_n\}) \leftrightarrow (\{a_n\})$ can be applicable to them, one will get the following equalities: $((\{a_1\}) \vee (\{a_2\})) = \top$ and $((\{a_1\}) \wedge (\{a_2\})) = \perp$.

Together these two equalities mean that after the measurement of the observable A – that is, in the moment when the records of the logical values of $(\{a_1\})$ and $(\{a_2\})$ are created – these propositions turn out to be *mutually exclusive*

$$(\{a_1, a_2\}) \equiv ((\{a_1\}) \oplus (\{a_2\})) = \top \quad . \quad (2.21)$$

Apropos of the normalization requirement $c_1^2 + c_2^2 = 1$, let us observe that it would match the operation of exclusive disjunction (2.21), if the squared norms c_1^2 and c_2^2 were viewed as the elements of the Boolean domain $\{\top, \perp\} \equiv \{1, 0\}$ corresponding respectively to the logical values

of the propositions $(\{a_1\})$ and $(\{a_2\})$ once these values were created in the measurement of A .

As regards the squared norms involved, according to Gleason's theorem (modified for the case of two-dimensional state spaces [26]), if one would like to assign a real valued function $m(\{a_n\}) \geq 0$ – e.g., a post-measurement logical value of the proposition $(\{a_n\})$, namely, $m(\{a_n\}) = (\{a_n\}) \in \{1, 0\}$ – to the vector $|a_n\rangle$ of the orthonormal set $\{|a_1\rangle, |a_2\rangle\}$ such that the following sum

$$\begin{aligned} m(\{a_1, a_2\}) &= m((\{a_1\}) \vee (\{a_2\})) \\ &= m(\{a_1\}) + m(\{a_2\}) = 1 \end{aligned} \quad (2.22)$$

holds true whenever the equality entailing the mutual exclusiveness of the outcome events $\{a_1\}$ and $\{a_2\}$

$$m((\{a_1\}) \wedge (\{a_2\})) = m(\{a_1\}) \times m(\{a_2\}) = 0 \quad (2.23)$$

is valid, then the only possible choice is $m(\{a_n\}) = |\langle a_n | \psi \rangle|^2$ in which $|\psi\rangle$ is an arbitrary but fixed vector $|\psi\rangle = c_1|a_1\rangle + c_2|a_2\rangle$.

For the sake of clarity and simplicity, let us suppose that $|\psi\rangle$ is the state of an *equal superposition* (i.e., a superposition in which all the coefficients have equal norms) such that this state can be written down as $|\psi\rangle = 1/\sqrt{2}(e^{i\phi_1}|a_1\rangle + e^{i\phi_2}|a_2\rangle)$ where phases ϕ_1 and ϕ_2 are real. As it can be readily seen, in that case the eigenstates $|a_1\rangle$ and $|a_2\rangle$ are interchangeable in $|\psi\rangle$ up to the phase factors of the superposition coefficients c_1 and c_2 , that is, $e^{i\phi_1}|a_1\rangle \leftrightarrow e^{i\phi_2}|a_2\rangle$.⁹

On the other hand, the exclusive disjunction $|c_1|^2 + |c_2|^2 = 1$ demonstrates that only the

⁸This evaluation is due to the easily verifiable assumption that the microsystem S can be prepared in two different eigenstates of the observable A and in a superposition of two such states.

⁹Such an interchangeability results by letting the unitary "swapping" operator $\hat{S} = e^{i\phi_1}|a_1\rangle\langle a_2|e^{-i\phi_2} + e^{i\phi_2}|a_2\rangle\langle a_1|e^{-i\phi_1}$ act on $|\psi\rangle$. This operator satisfies the invariance relation $\hat{S}|\psi\rangle = |\psi\rangle$.

magnitudes of the superposition coefficients c_1 and c_2 can correspond to the logical values $\{1, 0\}$ of the propositions of elementary events $(\{a_1\})$ and $(\{a_2\})$ once these values are created. That is, the phase factors $e^{i\phi_1}$ and $e^{i\phi_2}$ do not indicate which one of these propositions become 1. Accordingly, the symmetry $e^{i\phi_1}|a_1\rangle \leftrightarrow e^{i\phi_2}|a_2\rangle$ possessed by the equal superposition state $|\psi\rangle$ implies that prior to the measurement the propositions $(\{a_1\})$ and $(\{a_2\})$ are *indistinguishable from one another* except for their names, i.e., $(\{a_1\}) \leftrightarrow (\{a_2\})$.

So, considering that the equal superposition state $|\psi\rangle$ contains no physical evidence favoring one proposition over another, one can invoke *the principle of indifference* [15] and deduce from the a priori nonexistence of the logical values of $(\{a_1\})$ and $(\{a_2\})$ in the state $|\psi\rangle$ that both of these propositions are *equally likely* to become 1 during the measurement.¹⁰

In a word, due to the fact that the propositions of elementary events $(\{a_1\})$ and $(\{a_2\})$ are indistinguishable (equivalent) in the equal superposition state $|\psi\rangle$ and after the measurement correspond to the mutually exclusive events such that $(\{a_1\}) + (\{a_2\}) = 1$, one can assign both of these propositions *an equal probability* $\Pr(\{a_1\}) = \Pr(\{a_2\}) = 1/2$ of coming out 1 in the course of the measurement as $(\{a_1\})$ and $(\{a_2\})$ cannot be evaluated before the measurement.¹¹

This concludes the presentation of the way in which irreducible probabilities a central element of the Born rule appear in the intuitionistic interpretation of quantum mechanics in the

case of the superposition coefficients of equal magnitude and the two-dimensional state space of the measured system S .

The generalization of the presented account to the case of N coefficients with equal norms is rather straightforward¹². As to the extension of the account to the case of the superposition $|\psi\rangle = \sum_{n=1}^N c_n |a_n\rangle$ with the coefficients c_n of non-equal magnitudes, it can be achieved by means of logical theories of probability (see, for example, [28, 29]) which (in order to save the principle of indifference and extend it to the case of unsymmetrical evidence) assert that the elementary events $\{a_n\}$ may be assigned unequal *weights* w_n (that are nothing more than replication counts indicating duplicated observations) and probabilities can be computed whatever the evidence may be, symmetrical or not.

Along these lines, let us suppose that the squared norms of the coefficients c_n (of non-equal in general magnitudes) can be expressed as a fraction of two positive integers, namely, $|c_n|^2 = w_n / (\sum_{n=1}^N w_n)$. Then, the proposition of elementary event $(\{a_n\})$ can be brought into the following formal form

$$\begin{aligned} (\{a_n\}) &\equiv ((\{a_n\}_1) \vee (\{a_n\}_2) \cdots \vee (\{a_n\}_{w_n})) \\ &= \sum_{i=1}^{w_n} (\{a_n\}_i) \quad , \quad (2.24) \end{aligned}$$

where the subscript i indicates replicated disjoint events $\{a_n\}_i$ which are all identical with respect to $(\{a_n\})$. The association of the propositions $(\{a_n\})$ with the a priori probabilities $\Pr(\{a_n\}) =$

¹⁰This inference is in line with Jayne's invariance condition [27] according to which *equal probabilities should be assigned to equivalent propositions*.

¹¹The premeasurement logical values of the propositions $(\{a_n\})$ cannot be found (computed) by means of the Schrödinger equation and since *not-* (i.e., *other-than-*) Schrödinger dynamics is impossible, the probabilities of the events $\{a_n\}$ are interpreted as corresponding to a *genuine stochastic process*.

¹²Strictly speaking, the operation of exclusive disjunction, XOR, is true when an odd number of propositions $(\{a_n\})$ are true. So, in the case, in which the dimension N of the sample space of the measurement Ω is greater than 2, the proposition of a definite outcome must contain the additional to the operation XOR term B_N providing for the true output when only a single $(\{a_n\})$ is true. For example, when $N = 4$, this proposition takes the form $(\Omega) = (\text{XOR}_4 \wedge \neg B_4)$, where $\text{XOR}_4 = ((\{a_1\}) \oplus (\{a_2\}) \oplus (\{a_3\}) \oplus (\{a_4\}))$ and $B_4 = \max(\{(\{a_i\}) \times (\{a_j\}) \times (\{a_k\})\}_{i>j>k \in \{1,2,3,4\}})$.

$1/N$ has been already established in the special case where all the possible events $\{a_n\}$ have unity weights w_n . Thus, in the case of the weight $w_n > 1$, a similar line of reasoning can be used to prove the next formula

$$\Pr(\{a_n\}) = \sum_{i=1}^{w_n} \Pr(\{a_n\}_i) = \frac{w_n}{\sum_{n=1}^N w_n} . \quad (2.25)$$

So, the presented account is restricted to a finite dimensional sample space Ω and thus a finite dimensional state space. But even if a Hilbert space is countably infinite, the given account may still be held up whenever a replacement of a countably infinite orthonormal basis of the measured system S with some truncated (finite-dimensional) basis can be justified. *Such is the case of a typical microsystem* for which one can safely assume a discrete spectrum of energies limited by some finite upper level whose order of magnitude is similar to ones of energies of electrons in an atom or a solid.

However, the shown intuitionistic account may not be extended to an uncountable (i.e., uncountably infinite) Hilbert space *which is the case of a typical macroscopic system* whose degrees of freedom in particular the macroscopic ones can vary continuously and in an unconfined, unbounded manner. Apparently, this is equivalent to the elimination of the applicability of quantum probabilities and the Born rule to common macroscopic systems.

But then again, due to the equivalence of boundedness and continuity [30], unbounded linear Hermitian operators in an uncountable Hilbert space are discontinuous and thus generate *noncomputability* [31]. This means that the application of quantum formalism to a typical macroscopic system necessarily causes the existence of mathematical entities in the theoretical representation of the system that are incapable of being computed by any deterministic algorithm in any finite amount

of time.¹³ Clearly, those entities cannot be verifiable (demonstrable) which makes the quantum mechanical description of any typical macroscopic system constructively unprovable and for that reason inapplicable.

3 CONCLUSION

The essential question that may raise now is whether probabilities appearing in the intuitionistic interpretation are objective (i.e., fundamentally ingrained in nature) or subjective (i.e., due to the rational agent's ignorance). The peculiarity here is that neither answer to this question can be regarded as a correct one.

Really, according to the explanation presented in the previous section, quantum probabilities are introduced to account for the agent's lack of knowledge about the premeasurement logical values of the propositions of elementary events ($\{a_n\}$) in the case where the state $|\psi\rangle$ does not belong already (i.e., before the act of measurement) to one of the vectors $|a_n\rangle$ of the eigenbasis of the observable A that is going to be measured (and so the sample space of the measurement Ω contains more than one element). As long as these propositions are considered symmetrical with respect to the state $|\psi\rangle$, the agent has no other information than the number of the mutually exclusive events $\{a_n\}$ that can occur during the measurement. Consequently, the agent is justified in assigning each of the possible events the equal probability. In this manner, the resultant probabilities may be called *subjective* seeing as they reflect the agent's incomplete knowledge of the world.

But, on the other hand, quantum probabilities are *objective* in that they are a consequence of the fact that the decision problem of the Schrödinger equation $\Pi(H_C)$ is undecidable. That is, for this problem it is impossible to construct an algorithm (applicable to all systems, i.e., all allowable Hamiltonian operators H) that would always lead

¹³To be precise, one can no longer speak of the Hilbert space of a typical macroscopic system since "the observables of an infinite system usually have a host of physically inequivalent representations, corresponding to macroscopically different classes of states" [32].

to a correct 'true'-or-'false' answer. Hence, the origin of the undecidability of the problem $\Pi(H_C)$ is not subjective, i.e., it is not just a lack of agent's imagination: No matter how ingenious the rational agent is, there is no way to exactly solve the Schrödinger equation for any given physical system. As a result, the validity of the linear, deterministic Schrödinger evolution for an arbitrary system such as a typical macroscopic apparatus M cannot be proven constructively, i.e., by demonstrating a procedure of finding the solution set $\{|u\rangle\}$ of the Schrödinger equation for a completely arbitrary Hamiltonian operator.

The similar ambiguity characterizes the question whether or not the formalism of quantum mechanics in agreement with the intuitionistic interpretation can be considered complete. Definitely, as stated by the intuitionistic interpretation, the quantum description of reality can be considered *incomplete* as it contains the proposition $(\{a_1, a_2\}) = \{\}$ which is undecidable from within theory (the truth value of this proposition cannot be predicted through calculations with the Schrödinger equation), but which is, nonetheless, decidable through experiment (i.e., the measurement of the observable A). And yet, no further accumulation of information about the composite system $S+M$ could help the agent to decide (compute) the logical values of the propositions $(\{a_1\})$ and $(\{a_2\})$ before the measurement, inasmuch as the fact of the statement $(\{|u\rangle\} \neq \emptyset)$ being neither provable nor disprovable is not determined by any finite amount of the information available to the agent. In view of that fact, the presented intuitionistic account of quantum description may be regarded as *complete*.

COMPETING INTERESTS

Author has declared that no competing interests exist.

References

- [1] Tammaro E. Why Current Interpretations of Quantum Mechanics are Deficient. arXiv:1408.2093; 2014.
- [2] Deutsch D. Quantum theory of probability and decisions. Proceedings of the Royal Society of London, Series A. 1999;455:3129-3137.
- [3] Barnum H, Caves C, Finkelstein J, Fuchs C, Schack R. Quantum probability from decision theory? Proceedings of the Royal Society of London, Series A. 2000;456:1175-1182.
- [4] Wallace D. Everettian rationality: defending Deutsch's approach to probability in the Everett interpretation. Studies in History and Philosophy of Science, Part B. 2003;34(3):415-439.
- [5] Gill R. On an argument of David Deutsch. In M. Schurmann and U. Franz, editors, Quantum Probability and Infinite Dimensional Analysis: From Foundations to Applications. World Scientific, Singapore. 2005;18.
- [6] Assis A. On the Nature of $a_k^*a_k$ and the Emergence of the Born Rule. arXiv:1009.1532 [quant-ph]; 2012.
- [7] Eriksson L, Hájek A. What are Degrees of Belief? Studia Logica (Special Issue, Formal Epistemology). 2007;86(2):185-215.
- [8] Hájek A. Fifteen Arguments Against Hypothetical Frequentism. Erkenntnis. 2009;70:211-235.
- [9] Hemmo M, Pitowsky I. Quantum probability and many worlds. Studies in History and Philosophy of Modern Physics. 2007;38:333-350.
- [10] Zeh H. What is achieved by decoherence? arXiv:quant-ph/9610014; 1996.
- [11] Zeh H. There is no first quantization. Physics Letters A. 2003;309:329-334.
- [12] Zurek W. Probabilities from entanglement, Born's rule $p_k = |\psi_k|^2$ from envariance. Phys. Rev. A. 2005;71(5):052105.
- [13] Schlosshauer M, Fine A. On Zurek's derivation of the Born rule. Found. Phys. 2005;35(2):197-213.
- [14] Mohrhoff U. Probabilities from envariance? International Journal of Quantum Information. 2004;2(2):221-230.
- [15] Hájek A. Interpretations of Probability. In N. Zalta, editor, The Stanford Encyclopedia of Philosophy; 2012.

- [16] Easwaran K. The Varieties of Conditional Probability. In D. Gabbay, P. Bandyopadhyay, M. Forster, P. Thagard, and J. Woods, editors, *Philosophy of Statistics*. 2011;7:137-148.
- [17] Bolotin A. Intuitionistic interpretation of quantum mechanics. arXiv:1605.03125v1 [quant-ph]; 2016.
- [18] McGrath M. Propositions. In N. Zalta, editor, *The Stanford Encyclopedia of Philosophy*; 2014.
- [19] Caspers M, Heunen C, Landsman N, Spitters B. Intuitionistic quantum logic of an n-level system. *Foundations of Physics*. 2009;9(7):731-759.
- [20] Artemov S, Iemhoff R. The basic intuitionistic logic of proofs. *Journal of Symbolic Logic*. 2007;72:439-451.
- [21] Taranovsky D. Constructive Mathematical Truth. arXiv:math/0605138 [math.LO]; 2006.
- [22] Popelier P. Preface. In P. Popelier, editor, *Solving the Schroedinger Equation - Has Everything Been Tried?* Imperial College Press. 2011;XV-XIX.
- [23] Cubitt T, Perez-Garcia D, Wolf M. Undecidability of the spectral gap. *Nature*. 2015;528:207-211.
- [24] Chellas B. *Modal Logic: An Introduction*. Cambridge University Press, New York; 1980.
- [25] Leggett A. Testing the limits of quantum mechanics: motivation, state of play, prospects. *J. Phys.: Condens. Matter*. 2002;14:R415-R451.
- [26] De Zela F. Gleason-Type Theorem for Projective Measurements, Including Qubits: The Born Rule Beyond Quantum Physics. *Foundations of Physics*. 2016;1-14.
- [27] Jaynes E. *Probability Theory: The Logic of Science*. Cambridge University Press, New York; 2003.
- [28] Carnap R. *Logical foundations of probability*. University of Chicago Press, Chicago; 1950.
- [29] Vohanka V. Modality, logical probability, and the trinity: In defense of a weak skepticism. *Heusenstamm : Editiones Scholasticae*; 2014.
- [30] Hellman G. Constructive mathematics and quantum mechanics: Unbounded operators and the spectral theorem. *Journal of Philosophical Logic*. 1993;22:221-248.
- [31] Myrvold W. Computability in Quantum Mechanics. In W. Depauli-Schimanovich, E. Koehler, and F. Stadler, editors, *The Foundational Debate: Complexity and Constructivity in Mathematics and Physics*. Springer Netherlands, 1995;3:976-998.
- [32] Mundici D. Logic of Infinite Quantum Systems. *International Journal of Theoretical Physics*. 1993;32(10):1941-1955.

© 2017 Bolotin; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/18303>